



Carbene-Stabilized Dithiolene (L^0) Zwitterions

Yuzhong Wang, Phuong M. Tran, Yaoming Xie, Pingrong Wei, John N. Glushka, Henry F. Schaefer III, and Gregory H. Robinson*

Abstract: A series of reactions between Lewis bases and an imidazole-based dithione dimer (**1**) has been investigated. Both cyclic(alkyl)(amino)carbene (CAAC) (**2**) and *N*-heterocyclic carbene (NHC) (**4**), in addition to *N*-heterocyclic silylene (NHSi) (**6**), demonstrate the capability to cleave the sulphur-sulphur bonds in **1**, giving carbene-stabilized dithiolene (L^0) zwitterions (**3** and **5**) and a spirocyclic silicon-dithiolene compound (**7**), respectively. The bonding nature of **3**, **5**, and **7** are probed by both experimental and theoretical methods.

Largely due to the interplay of variable oxidation states of transition metals and the non-innocent character of the dithiolene ligand,^[1] the intriguing redox behavior of transition metal-based dithiolene complexes has fascinated chemists for more than a half-century.^[1,2] The oxidation state of the dithiolene ligand has previously been reported to vary between three ligand forms: dithiolate dianion (L^{2-}), radical monoanion ($L^\bullet-$), and neutral dithione (or dithiete) (L^0) (Figure 1).^[2,3]





Figure 1. Redox states of dithiolene ligands.

In contrast to the dominant roles of dithiolate (L^{2-}) ligands in transition metal dithiolene chemistry,^[2a] dithiolene ligands with monoanionic radical character ($L^\bullet-$) are receiving increasing attention.^[2f] For example, this laboratory recently reported the synthesis and molecular structure of lithium dithiolene radical (Scheme 1),^[4] which provides a unique synthetic platform for accessing the largely unexplored dithiolene chemistry of main group elements.^[5] In contrast to dithiolate dianions (L^{2-} , containing six π -electrons), the $4\pi\epsilon$ -containing neutral dithiolene (L^0) is an electron-deficient

How to cite: *Angew. Chem. Int. Ed.* **2021**, *60*, 22706–22710
International Edition: doi.org/10.1002/anie.202108498
German Edition: doi.org/10.1002/ange.202108498



Scheme 1. Synthesis of **3** ($R=2,6$ -diisopropylphenyl).

system, thus inclined to seek electron-rich species.^[2j] Neutral dithiolenes (L^0) have been reported to exist as dithiones and dithietes (Figure 1).^[2a,g,i,6] While structurally characterized dithiones and dithietes have been reported,^[6d,e,7] zwitterionic dithiolenes (L^0) (Figure 1) remain synthetic targets. Considering its electron deficiency, the dithiolene (L^0) system may employ one of its two sulphur atoms to accept one pair of electrons from a Lewis base ligand. This would result in charge separation, giving a Lewis base-stabilized dithiolene (L^0) zwitterion. To this end, we report the syntheses,^[8] structures^[8] and computations^[8] of the first carbene-stabilized neutral dithiolene (L^0) zwitterions (**3** and **5**). Notably, when employing NHSi as the Lewis base, a spirocyclic silicon-dithiolene compound (**7**) was achieved.

Scission of sulphur-sulphur bonds is important in biological processes and organic synthesis,^[9] which usually involves transition metals or main group species (i.e., nucleophiles, electrophiles, and FLPs).^[9,10] Carbene-insertion into the S–S bond is attracting increasing attention.^[11] In particular, transient donor/acceptor-free carbenes have been shown to cleave the S–S bond through ylide formation or radical mechanisms.^[11d] This laboratory recently synthesized the imidazole-based dithione dimer, containing S–S bonds, (**1**) by one-electron oxidation of the corresponding lithium dithiolene radical anion (Scheme 1).^[12] The analogues of **1** have been synthesized by both sulfurization of imidazole-based dione with Lawesson's reagent and Br_2 oxidation of nickel bis-dithiolene complex.^[13] Dissociation of these dithione dimers in solution has not been observed. Dithione was only documented as a reactive intermediate in the synthesis of stable analogue of **1** ($R=iPr$).^[13a] The 1:2 reaction of **1** with $^{Me}CAAC$ (**2**)^[14] in toluene results in the cleavage of the S–S bonds in **1**, affording the carbene-

[*] Dr. Y. Wang, P. M. Tran, Dr. Y. Xie, Dr. P. Wei, Dr. J. N. Glushka, Prof. Dr. H. F. Schaefer III, Prof. Dr. G. H. Robinson
Department of Chemistry, Center for Computational Chemistry, and Complex Carbohydrate Research Center, The University of Georgia Athens, GA 30602-2556 (USA)
E-mail: robinson@uga.edu

Supporting information and the ORCID identification number(s) for the author(s) of this article can be found under:
<https://doi.org/10.1002/anie.202108498>.

stabilized dithiolene (L^0) zwitterion (**3**) as a purple-blue powder in 94.0% yield (Scheme 1).^[8] X-ray single crystals of **3** were obtained by recrystallization from hot toluene (or from a 1,2-difluorobenzene/hexane mixed solvent). The UV–Vis absorption spectrum of **3** in toluene reveals one strong broad absorption ($\lambda_{\text{max}} = 611 \text{ nm}$) and one weak broad absorption ($\lambda_{\text{max}} = 451 \text{ nm}$) (Figure S1).^[8] Due to poor solubility in benzene (or toluene) at room temperature, the ^{13}C NMR spectrum of **3** (in C_6D_6) was determined at an elevated temperature ($T = 68^\circ\text{C}$).^[8] The ^{13}C NMR resonance of the carbene carbon in **3** [C(28) in Figure 2] (in C_6D_6) is observed at 208.53 ppm, which shifts upfield compared to that (313.6 ppm) of the free ${}^{\text{Me}}\text{CAAC}$ ligand (in C_6D_6).^[14a]

X-ray structural analysis^[8] reveals that **3** exists as a pair of enantiomers (with identical bonding parameters) (Figure 2), which may be attributed to the steric congestion of **3**, thereby restricting the rotation about the C(2)–S(2) bond (see the space filling model of **3** in Figure S2).^[8] In addition, the two 2,6-diisopropylphenyl substituents of the imidazole ring are bent toward the opposite sides of the five-membered ring. While being marginally shorter than the corresponding theoretical values in **3** model ($d_{\text{C-C}} = 1.393 \text{ \AA}$, $d_{\text{C-S}} = 1.704$ and 1.735 \AA),^[8] the C_2S_2 unit in **3** shows an elongated C–C bond [$1.374(2) \text{ \AA}$] and concomitantly shortened C–S bonds

[$1.6941(16)$ and $1.7137(15) \text{ \AA}$] compared to those in **1** [$d_{\text{C-C}} = 1.362(8) \text{ \AA}$, $d_{\text{C-S}} = 1.724(6)$ and $1.733(6) \text{ \AA}$.^[12] By comparison, the C–S bonds in the C_2S_2 unit of **3** [$1.6941(16)$ and $1.7137(15) \text{ \AA}$] are longer than the C(1)=S(1) double bond of **3** [$1.6718(16) \text{ \AA}$], however, shorter than the typical C–S single bonds (such as those in $\text{PtPh}_2(9\text{S}3)$ [$9\text{S}3 = 1,4,7\text{-trithiacyclo-}9\text{~methyl-}5\text{-octene}$] [$1.815(8)$ – $1.848(8) \text{ \AA}$]).^[15] The C(3)-C(2)-S(2)-C(28) torsion angle (-63.7°) of **3** is comparable to that (-63.1°) in **3** model, whereas the S(2)-C(2)-C(3)-S(3) torsion angle (9.0°) of **3** is somewhat smaller than its theoretical value (16.5°) (which may be due to packing effects).^[16] The carbene carbon in **3** [i.e., C(28)] exhibits a trigonal planar geometry [the sum of bond angles around C(28) = 359.7°].

The DFT computations^[8] of **3** at the B3LYP/6-311G** level show that the HOMO is predominantly dithiolene-based, involving polarized C–C π -bonding and C–S π -antibonding character, whereas the LUMO is mainly CAAC-based with $\text{C}_{\text{CAAC}}=\text{N}$ π -antibonding character (Figure 3). NBO natural population analysis reveals a positive charge of +0.43 at the S(2) atom and a negative charge of -0.35 at the S(3) atom of **3**, supporting **3A** (Figure 4) as the major resonance contributor. It is reasonable to deduce that formation of **3** may be through CAAC-mediated heterolytic cleavage of S–S bonds in **1**. The diradical product **3**[·] would be expected if CAAC-mediated cleavage of S–S bonds in **1** is homolytic and involves a radical mechanism. However, our computations reveal that the triplet diradical **3**[·] is 15.32 kcal mol⁻¹ higher in energy than the singlet zwitterionic **3**.^[8]

We also investigated the corresponding reactions of **1** with other Lewis bases (i.e., NHC and NHSi). The 1:2 reaction of **1** with $\text{NHC}^{\text{iPr}}[17]$ (**4**) gave enantiomeric **5** (the NHC analogue of **3**) as a yellow powder (in 84.4% yield) (Scheme 2). Unexpectedly, the parallel reaction of **1** with sterically demanding $\text{NHSi}^{\text{Dipp}}[18]$ (Dipp = 2,6-diisopropylphenyl) (**6**)




Figure 2. Molecular structures of the enantiomers of **3**. Thermal ellipsoids represent 30% probability. Hydrogen atoms have been omitted for clarity. Selected bond distances (\AA) and angles (deg): S(1)–C(1), 1.6718(16); C(2)–C(3), 1.374(2); C(2)–S(2), 1.7137(15); C(3)–S(3), 1.6941(16); S(2)–C(28), 1.7285(16); C(28)–C(31), 1.524(2); C(28)–N(3), 1.304(2); C(2)–C(3)–S(3), 131.42(12); C(2)–S(2)–C(28), 106.69(8).




Figure 3. Frontier molecular orbitals of **3**.







Figure 4. Canonical structures of **3** and **3**[·].




quantitatively resulted in colourless spirocyclic silicon–dithiolene complex (**7**) (Scheme 2).^[8] Consequently, **7** joins a small group of structurally characterized silicon–dithiolene complexes.^[5c,19] To the best of our knowledge, the literature does not reveal a report of stable carbene-, or silylene-, mediated cleavage of a sulphur–sulphur bond (in a molecular compound). The dramatic colour difference between **3** and **5** may be related to their distinct HOMO–LUMO energy gaps [$\Delta E_{\text{HOMO-LUMO}} = 2.08 \text{ eV}$ (**3**) and 2.88 eV (**5**)].^[8] Compound **5** is largely insoluble in benzene and toluene. The ^1H NMR spectrum of **5** (Figure S6) in $\text{C}_6\text{D}_5\text{Br}$ at room temperature shows broad lines. Notably, the methine proton resonance of the NHC^{iPr} ligand in **5** cannot be even distinguished from the baseline. Variable-temperature (VT) ^1H NMR experiments of **5** (Figure S5) show that with the temperature increasing from 25 to 80°C , the methyl proton resonances at 0.74 and 1.04 ppm evolve from broad singlets to sharp doublets. Meanwhile, the methine proton resonance of the NHC^{iPr} unit of **5** is observed as a broad singlet at 5.68 ppm (Figure S7). Such dynamic behaviour of **5** in $\text{C}_6\text{D}_5\text{Br}$ should be due to steric congestion of this compound, which restricts the ligand rotation (see the space filling model of **5** in Figure S2). In contrast to carbene-complexed zwitterions (**3** and **5**), spirocyclic compound **7** is readily soluble in toluene. The four-coordinate silicon atom in **7** exhibits a ^{29}Si NMR resonance at -4.67 ppm . By comparison, the ^{29}Si NMR resonance of the silicon bis(dithiolene) complex, $\text{Si}[\text{S}_2(\text{o-C}_6\text{H}_4)]_2$,^[19a] is observed at 45.4 ppm , while those for the alkoxy-functionalized NHSi sulphides were observed at -24.3 and -24.4 ppm .^[20] Spirocyclic silanes have been obtained via 4 + 1 cyclization reactions of silylenes with various dienes.^[21] Thus, **7**, may also be accessed through 1,2-dithiones (Figure 1).

Similar to **3**, the NHC-complexed dithiolene (L^0) zwitterion (**5**) also exists as an enantiomeric pair (with identical bonding parameters) (Figure 5a).^[8] By comparison with **3**, the most obvious structural changes for **5** are the $\text{C}(3)\text{-C}(2)\text{-S}(2)$ –



(a)



(b)

Figure 5. Molecular structures of enantiomeric **5** (a) and spiro silicon–dithiolene **7** (b). Thermal ellipsoids represent 30% probability. Hydrogen atoms have been omitted for clarity. Selected bond distances (\AA) and angles (deg): For **5**, $\text{S}(1)\text{-C}(1)$, $1.673(2)$; $\text{C}(2)\text{-C}(3)$, $1.380(3)$; $\text{C}(2)\text{-S}(2)$, $1.718(2)$; $\text{C}(3)\text{-S}(3)$, $1.690(2)$; $\text{S}(2)\text{-C}(28)$, $1.757(2)$; $\text{C}(2)\text{-C}(3)\text{-S}(3)$, $131.22(17)$; $\text{C}(3)\text{-C}(2)\text{-S}(2)$, $125.84(17)$; $\text{C}(2)\text{-S}(2)\text{-C}(28)$, $106.62(11)$. For **7**, $\text{S}(1)\text{-C}(1)$, $1.6638(9)$; $\text{C}(2)\text{-C}(3)$, $1.3468(12)$; $\text{C}(2)\text{-S}(2)$, $1.7374(9)$; $\text{C}(3)\text{-S}(3)$, $1.7339(9)$; $\text{S}(2)\text{-Si}(1)$, $2.1591(4)$; $\text{N}(3)\text{-Si}(1)$, $1.7118(9)$; $\text{C}(2)\text{-C}(3)\text{-S}(3)$, $125.12(7)$; $\text{C}(2)\text{-S}(2)\text{-Si}(1)$, $94.73(3)$; $\text{S}(2)\text{-Si}(1)\text{-S}(3)$, $100.893(15)$.

$\text{C}(28)$ torsion angle (-63.7° for **3**, -114.8° for **5**) and $\text{N}(3)\text{-C}(28)\text{-S}(2)\text{-C}(2)$ torsion angle ($+154.1^\circ$ for **3**, -126.5° for **5**). These structural distinctions between **3** and **5** are largely due

to the different steric effects of the carbene ligands (**2** and **4**). The +0.33 positive charge of the S(2) atom and -0.35 negative charge of the S(3) atom in **5** compare well to that in **3** [+0.43 for S(2) and -0.35 for S(3)], suggesting its zwitterionic character. The 1.34–1.48 Wiberg bond index of the C–S_{terminal} bonds [i.e., C(1)–S(1) and C(3)–S(3) bonds] in **3** and **5** indicate that these two bonds have modest double bond character. The C_{carbene}–S bond in **5** [1.757(2) Å; 57.5 % toward C and 42.5 % toward S] is somewhat longer and more polarized than that in **3** [1.7285(16) Å; 55.3 % toward C and 44.7 % toward S]. Frontier molecular orbitals of **5** (Figure S12) are comparable with those of **3**.^[8]

X-ray structural analysis^[8] (Figure 5b) shows that the four-coordinate silicon atom in **7** adopts a distorted tetrahedral geometry ($\tau = 0.78$).^[22] Similar to that [2.1381(6) Å] of Si[S₂(*o*-C₆H₄)₂] containing a four-coordinate silicon atom,^[19a] the Si–S bond distances of **7** [2.1604(4) Å, av] are much shorter than those [2.3195(12)–2.3374(12) Å] in the reported silicon tris(dithiolene) dianion containing a six-coordinate silicon atom.^[5c] The bonding parameters of the C₂S₂ unit [$d_{C-S} = 1.7357(9)$ Å, av; $d_{C=C} = 1.3468(12)$ Å] of **7** are consistent with those for the reported dithiolates.^[2a,23] Thus, the Si^{II} atom in NHSi^{DiPP} (**6**) is oxidized to Si^{IV} when forming **7**. The natural charge residing at the central silicon of **7** is +1.59. While the HOMO of **7** is dithiolene-based (containing both C–C π-bonding and C–S π-antibonding character), the LUMO involves silicon–sulphur σ-antibonding interactions (Figure S13).^[8]

The sulphur–sulphur bonds in the imidazole-based dithione dimer (**1**) have successfully been cleaved using a series of stable Lewis bases [³MeCAAC (**2**), NHC^{iPr} (**4**), and NHSi^{DiPP} (**6**)], which gives carbene-stabilized dithiolene (L⁰) zwitterions (**3** and **5**) and spirocyclic silicon–dithiolene complex (**7**), respectively. The intriguing redox reactivity of these compounds are being further investigated in this laboratory.

Acknowledgements

We are grateful to the National Science Foundation for support: CHE-1855641 (G.H.R., Y.W.) and CHE-2134792 (H.F.S.).

Conflict of Interest

The authors declare no conflict of interest.

Keywords: carbenes · dithiolenes · silicon · silylenes · sulphur

- [1] R. Eisenberg, H. B. Gray, *Inorg. Chem.* **2011**, *50*, 9741–9751.
- [2] a) *Dithiolene Chemistry: Synthesis Properties, and Applications* (Ed.: E. I. Stiefel), Wiley, Hoboken, NJ, **2004**; b) R. Kato, *Chem. Rev.* **2004**, *104*, 5319–5346; c) F. J. Hine, A. J. Taylor, C. D. Garner, *Coord. Chem. Rev.* **2010**, *254*, 1570–1579; d) S. Rabaca, M. Almeida, *Coord. Chem. Rev.* **2010**, *254*, 1493–1508; e) B. Garreau-de Bonneval, K. I. M. C. Ching, F. Alary, T. T. Bui, L. Valade, *Coord. Chem. Rev.* **2010**, *254*, 1457–1467; f) S. Sproules, K. Wieghardt, *Coord. Chem. Rev.* **2011**, *255*, 837–860; g) S.

- Sproules, *Prog. Inorg. Chem.* **2014**, *58*, 1–144; h) A. Kobayashi, E. Fujiwara, H. Kobayashi, *Chem. Rev.* **2004**, *104*, 5243–5264; i) P. Basu, K. J. Colston, B. Mogesa, *Coord. Chem. Rev.* **2020**, *409*, 213211; j) G. N. Schrauzer, V. Mayweg, *J. Am. Chem. Soc.* **1962**, *84*, 3221–3221; k) H. B. Gray, E. Billig, R. Williams, I. Bernal, *J. Am. Chem. Soc.* **1962**, *84*, 3596–3597; l) H. B. Gray, E. Billig, *J. Am. Chem. Soc.* **1963**, *85*, 2019–2020; m) A. Davison, R. H. Holm, N. Edelstein, A. H. Maki, *Inorg. Chem.* **1963**, *2*, 1227–1232.
- [3] B. S. Lim, D. V. Fomitchev, R. H. Holm, *Inorg. Chem.* **2001**, *40*, 4257–4262.
- [4] Y. Wang, H. P. Hickox, Y. Xie, P. Wei, S. A. Blair, M. K. Johnson, H. F. Schaefer III, G. H. Robinson, *J. Am. Chem. Soc.* **2017**, *139*, 6859–6862.
- [5] a) Y. Wang, Y. Xie, P. Wei, S. A. Blair, D. Cui, M. K. Johnson, H. F. Schaefer III, G. H. Robinson, *Angew. Chem. Int. Ed.* **2018**, *57*, 7865–7868; *Angew. Chem.* **2018**, *130*, 7991–7994; b) Y. Wang, N. A. Maxi, Y. Xie, P. Wei, H. F. Schaefer III, G. H. Robinson, *Chem. Commun.* **2019**, *55*, 8087–8089; c) Y. Wang, C. A. Tope, Y. Xie, P. Wei, J. L. Urbauer, H. F. Schaefer III, G. H. Robinson, *Angew. Chem. Int. Ed.* **2020**, *59*, 8864–8867; *Angew. Chem.* **2020**, *132*, 8949–8952; d) P. M. Tran, Y. Wang, Y. Xie, P. Wei, H. F. Schaefer III, G. H. Robinson, *Chem. Commun.* **2021**, *57*, 2543–2546.
- [6] a) W. Kusters, P. D. Mayo, *J. Am. Chem. Soc.* **1973**, *95*, 2383–2384; b) E. A. M. Geary, L. J. Yellowlees, S. Parsons, L. Pilia, A. Serpe, M. L. Mercuri, P. Deplano, S. J. Clark, N. Robertson, *Dalton Trans.* **2007**, 5453–5459; c) M. R. Green, N. Jubran, B. E. Bursten, D. H. Busch, *Inorg. Chem.* **1987**, *26*, 2326–2332; d) P. C. Servaas, D. J. Stukens, A. Oskam, P. Vernooyjs, E. J. Baerends, D. J. A. Deridder, C. H. Stam, *Inorg. Chem.* **1989**, *28*, 4104–4113; e) T. Shimizu, H. Murakami, Y. Kobayashi, K. Iwata, N. Kamigata, *J. Org. Chem.* **1998**, *63*, 8192–8199.
- [7] M. G. B. Drew, J. M. Kisenyi, G. R. Willey, *J. Chem. Soc. Dalton Trans.* **1982**, 1729–1732.
- [8] See the supporting information for synthetic, computational, and crystallographic details. Deposition Numbers 2097537 (**3**), 2097538 (**5**), and 2097539 (**7**) contain the supplementary crystallographic data for this paper. These data are provided free of charge by the joint Cambridge Crystallographic Data Centre and Fachinformationszentrum Karlsruhe Access Structures service.
- [9] a) A. J. Parker, N. Kharasch, *Chem. Rev.* **1959**, *59*, 583–628; b) J. L. Kice, *Acc. Chem. Res.* **1968**, *1*, 58–64; c) D. N. Harpp, R. A. Smith, *J. Am. Chem. Soc.* **1982**, *104*, 6045–6053.
- [10] M. A. Dureen, G. C. Welch, T. M. Gilbert, D. W. Stephan, *Inorg. Chem.* **2009**, *48*, 9910–9917.
- [11] a) X. Hong, L. Lu, Q. L. Shen, *Synlett* **2019**, *30*, 1602–1606; b) Y. Kawamura, K. Akitomo, M. Oe, T. Horie, M. Tsukayama, *Tetrahedron Lett.* **1997**, *38*, 8989–8992; c) M. Hamaguchi, T. Misumi, T. Oshima, *Tetrahedron Lett.* **1998**, *39*, 7113–7116; d) Y. Zheng, R. J. Bian, X. L. Zhang, R. W. Yao, L. H. Qiu, X. G. Bao, X. F. Xu, *Eur. J. Org. Chem.* **2016**, 3872–3877; e) C. Q. Rao, S. Y. Mai, Q. L. Song, *Chem. Commun.* **2018**, *54*, 5964–5967; f) H. R. Yuan, T. Nuligonda, H. Y. Gao, C. H. Tung, Z. H. Xu, *Org. Chem. Front.* **2018**, *5*, 1371–1374.
- [12] Y. Wang, Y. Xie, P. Wei, H. F. Schaefer III, G. H. Robinson, *Dalton Trans.* **2019**, *48*, 3543–3546.
- [13] a) F. Bigoli, P. Deplano, F. A. Devillanova, J. R. Ferraro, V. Lippolis, P. J. Lukes, M. L. Mercuri, M. A. Pellinghelli, E. F. Trogu, J. M. Williams, *Inorg. Chem.* **1997**, *36*, 1218–1226; b) P. Deplano, M. L. Mercuri, G. Pintus, E. F. Trogu, *Comments Inorg. Chem.* **2001**, *22*, 353–374; c) F. Bigoli, S. Currelli, P. Deplano, L. Leoni, M. L. Mercuri, M. A. Pellinghelli, A. Serpe, E. F. Trogu, *J. Chem. Soc. Dalton Trans.* **2002**, 1985–1991.
- [14] a) C. Müller, D. M. Andrade, I. A. Bischoff, M. Zimmer, V. Huch, N. Steinbrück, A. Schäfer, *Organometallics* **2019**, *38*,

- 1052–1061; b) M. Soleilhavoup, G. Bertrand, *Acc. Chem. Res.* **2015**, *48*, 256–266.
- [15] M. A. Bennett, A. J. Canty, J. K. Felixberger, L. M. Rendina, C. Sunderland, A. C. Willis, *Inorg. Chem.* **1993**, *32*, 1951–1958.
- [16] J. W. Steed, in *Frontiers in Crystal Engineering*, 1st ed. (Eds.: E. R. T. Tiekkink, J. Vittal), Wiley, Chichester, UK, **2006**, pp. 67–90.
- [17] N. Kuhn, T. Kratz, *Synthesis* **1993**, *561*–562.
- [18] a) L. B. Kong, J. Y. Zhang, H. B. Song, C. M. Cui, *Dalton Trans.* **2009**, *5444*–*5446*; b) P. Zark, A. Schäfer, A. Mitra, D. Haase, W. Saak, R. West, T. Müller, *J. Organomet. Chem.* **2010**, *695*, 398–408.
- [19] a) U. Herzog, U. Böhme, G. Rheinwald, *J. Organomet. Chem.* **2000**, *612*, 133–140; b) R. Tacke, M. Mallak, R. Willeke, *Angew. Chem. Int. Ed.* **2001**, *40*, 2339–2341; *Angew. Chem.* **2001**, *113*, 2401–2403; c) O. Seiler, M. Buttner, M. Penka, R. Tacke, *Organometallics* **2005**, *24*, 6059–6062; d) K. Junold, C. Burschka, R. Bertermann, R. Tacke, *Dalton Trans.* **2010**, *39*, 9401–9413.
- [20] Y. Wang, Y. Xie, P. Wei, S. A. Blair, D. Cui, M. K. Johnson, H. F. Schaefer III, G. H. Robinson, *J. Am. Chem. Soc.* **2020**, *142*, 17301–17305.
- [21] M. Asay, C. Jones, M. Driess, *Chem. Rev.* **2011**, *111*, 354–396.
- [22] L. Yang, D. R. Powell, R. P. Houser, *Dalton Trans.* **2007**, *955*–964.
- [23] J. G. Breitzer, A. I. Smirnov, L. F. Szczepura, S. R. Wilson, T. B. Rauchfuss, *Inorg. Chem.* **2001**, *40*, 1421–1429.

Manuscript received: June 25, 2021

Revised manuscript received: July 23, 2021

Accepted manuscript online: July 27, 2021

Version of record online: September 12, 2021