

# Counterintuitive Chemistry: Carbene Stabilization of Zero-Oxidation State Main Group Species

Yuzhong Wang and Gregory H. Robinson\*



Cite This: *J. Am. Chem. Soc.* 2023, 145, 5592–5612



Read Online

ACCESS |


Metrics & More

Article Recommendations

**ABSTRACT:** Carbenes have evolved from transient laboratory curiosities to a robust, diverse, and surprisingly impactful ligand class. A variety of different carbenes have significantly contributed to the development of low-oxidation state main group chemistry. This Perspective focuses upon advances in the chemistry of carbene complexes containing main group element cores in the formal oxidation state of zero, including their diverse synthetic strategies, unusual bonding and structural motifs, and utility in transition metal coordination chemistry and activation of small molecules.

## INTRODUCTION

Carbenes, neutral molecules possessing a divalent carbon atom with six valence electrons, have evolved from transient laboratory curiosities to a robust, diverse, and surprisingly impactful ligand class. The first stable carbene,  $\lambda^3$ -phosphino-carbene, was reported by Bertrand in 1988.<sup>1,2</sup> Although reports of N-heterocyclic carbene (NHC, Figure 1)-based transition



**Figure 1.** Representative divalent carbon(II) (NHC and CAAC) and divalent carbon(0) compounds (i.e., carbenes) (CDP and CDC) and resonance structures of CDP and CDC: carbene (A) and allene (B).

metal complexes date back to the 1960s,<sup>3</sup> Arduengo reported the first stable N-heterocyclic carbene only three decades ago.<sup>4</sup> These seminal discoveries were prelude to a frenetic period of unprecedented advances in the development of carbene chemistry. Perhaps expectedly, the variety of stable carbenes has expanded dramatically. Notably, *nonclassical* carbenes, species containing one, or even no heteroatom in positions  $\alpha$  to the carbene carbon center, are presently receiving increased attention.<sup>5</sup> Of particular note are cyclic (alkyl)(amino)-carbenes (CAACs, Figure 1),<sup>6–8</sup> first reported by Bertrand in 2005.<sup>9</sup> CAACs exhibit electronic properties that are quite distinct from classical N-heterocyclic carbenes. Notably, both

computational and experimental data suggest that CAACs are not only stronger  $\sigma$ -donors but also more potent  $\pi$ -acceptors than NHCs.<sup>7,10–13</sup> Although carbenes continue to be extensively utilized throughout the whole of chemistry, their impact in organic synthesis, catalysis,<sup>14</sup> and the development of low-oxidation-state main group chemistry has been particularly significant.<sup>6,7,15–19</sup>

As evidenced by such iconic molecules as Ni(CO)<sub>4</sub> and (C<sub>6</sub>H<sub>6</sub>)<sub>2</sub>Cr, transition metals can readily adopt the formal oxidation state of zero. However, embracing the formal oxidation state of zero is a considerably less common endeavor for main group elements. The stabilization of molecules containing main group elements in the zero-oxidation state has become a popular research area over the past two decades.<sup>16,20–24</sup> To this end, carbenes—both NHCs and CAACs—are particularly attractive, considering their strong electron-donating capabilities coupled with their tunable electronic and steric properties.

This Perspective will chronologically review seminal advances in the counterintuitive chemistry of carbene-stabilized zero-oxidation state main group species—logically proceeding from atomic moieties to more complicated allotropic entities.

## CARBENE-STABILIZED E<sub>1</sub>(0) SPECIES

While diamond and graphite—the two most common carbon allotropes—possess remarkable industrial utility, the carbon(0) atom, the simplest building block of these stable carbon allotropes, is highly reactive. As a result, the study of complexes containing a carbon(0) core is intriguing. The first compounds

Published: March 6, 2023




containing a carbon(0) core are traced to the carbodiphosphorane (CDP) (Figure 1),  $(\text{PPh}_3)_2\text{C}$ , synthesized by Ramirez in 1961.<sup>25</sup> Decades later Frenking further clarified the “carbon(0) compound” essence of CDPs with theoretical and experimental data.<sup>26</sup> Theoretical evidence for the presence of a divalent carbon(0) core in  $(\text{NHC})_2\text{C}$  complexes (i.e., carbodicarbenes, CDCs, Figure 1) was also obtained.<sup>27,28</sup>

Consistent with the molecular orbitals of CDPs, the HOMO and HOMO-1 of CDCs correspond to the  $\pi$ -type and  $\sigma$ -type lone pair orbitals, respectively.<sup>27</sup> For CDPs and CDCs, while the  $\sigma$ -type lone pair orbital is localized at the central carbon atom, the  $\pi$ -type lone pair orbital largely resides at the central carbon possessing a measure of delocalization over the phosphine or carbene ligands. The bonding between the central C(0) atom and the ligands in CDPs and CDCs, therefore, can be best described as donor–acceptor interactions (resonance structure A, Figure 1). Singlet carbenes (such as NHCs and CAACs) have a lone pair of electrons and an empty p orbital on the carbene carbon atom (Figure 1). Consequently, the most remarkable distinction between carbenes and carbones is the unique double donor capability of the former.

Soon after the theoretical prediction by Frenking,<sup>27</sup> Bertrand synthesized the first carbodicarbene (**3**) from bis(*N*-methylbenzimidazol-2-yl)methane (**1**) (Scheme 1) in 2008.<sup>29</sup> X-ray

### Scheme 1. Synthesis of Carbodicarbene 3



structural analysis shows that the two N–C1–N planes in **3** are twisted by  $69^\circ$  (Figure 2). While the C1–C2 bond distance




Figure 2. Molecular structure of **3**.


of **3** [1.343(2) Å] compares well to the theoretical value (1.355 Å),<sup>30</sup> the C2–C1–C2' angle of **3** [ $134.8(2)^\circ$ ] is more acute than the calculated data ( $142.5^\circ$ ). This difference may be attributed to the intermolecular interactions in crystals and the shallow bending potential of carbon(0) compounds.<sup>30</sup>

Tetraaminoallenes are powerful nucleophiles and bases. Computations support the  $(\text{R}_2\text{N})_2\text{C} \rightarrow \text{C}=\text{C}(\text{NR}_2)_2$  bonding description, wherein the central carbon atom bears two lone pairs of electrons.<sup>27</sup> Both carbodicarbene **3**<sup>29</sup> and the tetraaminoallene  $[(\text{Me}_2\text{N})_2\text{C}]_2\text{C}$ <sup>31</sup> were reported to form an  $\eta^1$  complex with transition metal species, whereas reactions of “regular allenes” with transition metal species give  $\eta^2$  complexes involving one of the two  $\text{C}=\text{C}$   $\pi$  bonds.<sup>32</sup> While

carbenes are widely recognized as single  $\sigma$ -donors, carbones, bearing two electron pairs at the central carbon(0) centers, may serve as double donors. Although dimetalations of the carbene centers of CDPs are well documented,<sup>33</sup> synthesis of CDC-based geminal-bimetallic complexes is still challenging. The dormancy of the  $\pi$ -type lone pair of the C(0) center of CDC may be ascribed to back-donation of electron density from the carbon(0) atom to the NHC moieties. Ong recently made a breakthrough in this field by suppressing the  $\pi$ -acidity of the flanked NHC moiety in CDC.<sup>34</sup>


Geminal-bimetallic CDC complexes **5**, **6**, and **8** were synthesized using the corresponding CDC-based mononuclear complexes (**4** and **7**) (Scheme 2).<sup>34</sup> Both X-ray structural

### Scheme 2. Synthesis of Gem-Double Dative Bimetallic CDC Complexes (**5**, **6**, and **8**)



analyses and theoretical studies of **5**, **6**, and **8** support that the carbene center of the cyclometalated CDC moiety provides two lone electron pairs to the two metal centers via two dative bonds.

The chelating binding of CDC, via C–H bond activation, has twisted the flanking NHC moiety in these geminal-bimetallics of CDC, thereby decreasing the  $\pi$ -conjugation within the allenic framework (Figure 3). The solid-state structural analysis also revealed the presence of noncovalent ligand–ligand interactions (Figure 3) [between acetate (L:) and CDC for **5** and **6** and between bromide (L:) and CDC for **8**]. Thus, Ong suggested that the electron density from the




**Figure 3.** Activation of the second lone pair of CDC via chelating effect and ligand–ligand interaction.

external ligand (L:) (i.e., acetate or bromide) may be donated to the flanked NHC moiety of CDC and thus quench its  $\pi$ -acidity. As a result, the second dormant lone pair of the carbon(0) center may be activated via the cooperative effect between the chelating mode of CDC and ligand–ligand interactions. Recent observation of 1,2-addition of E–H bonds (E = B, C, Si) across the CDC central carbon atom and that of the flanking NHC unit in CDC, coupled with the results of X-ray structural analysis, indicates significant  $\pi$ -acidity of the  $C_{\text{NHC}}$  atom in CDC.<sup>35</sup> This finding is unexpected since NHCs are well-known as strong  $\sigma$ -donors, but weak  $\pi$ -acceptors. Thus, CDCs, bearing both an electron-rich carbene center and two  $\pi$ -acidic  $C_{\text{NHC}}$  centers, are expected to exhibit frustrated Lewis pairs (FLPs)-like reactivity.<sup>35</sup> Consequently, potential applications of such CDCs in catalysis and small molecule activation are promising.<sup>36,37</sup>

In addition to offering double-donor functionality, CDCs have also exhibited reactivity quite distinct from carbenes. While reaction of the bulky  $\text{NHC}^{\text{Dipp}}$  ligand (**9**) with  $\text{BH}_3 \cdot \text{THF}$  gives the 1:1 adduct (**10**),<sup>38</sup> reaction of CDC (**11**) with  $\text{BH}_3 \cdot \text{THF}$  gives an unusual three-coordinate dicationic hydrido boron complex (**12**) (Scheme 3).<sup>39</sup> Notably, computations

### Scheme 3. Synthesis of **10** and **12**




indicate that the C(0)-to-B electron donation involves two electron pairs of the CDC ligand (**11**). Synthesis of the dicationic borane complex via CDC coordination reveals the possible broad utility of CDCs in stabilizing highly charged main group species. The same CDC ligand (**11**) was subsequently employed in the synthesis of CDC-based di- and tricationic bismalkenes, involving C(0)-to-Bi double dative bonds.<sup>40</sup> Redox noninnocent ligands have attracted increased attention in the field of catalysis. England recently reported the redox noninnocence of CDC ligands in highly oxidized chromium and cobalt complexes.<sup>41</sup>

Theoretical studies<sup>42</sup> suggest that the *bent* trisilaallene, reported by Kira et al. in 2003, may also be interpreted as a


silylone ( $\text{L}_2\text{Si}$ , where L = a cyclic silylene ligand).<sup>43</sup> The first carbene-based silylone (**14**) was obtained by Roesky in 2013 via potassium graphite reduction of  $^{\text{Me}}\text{CAAC}$ -complexed  $\text{SiCl}_2$  (**13**) (Scheme 4).<sup>44</sup> **13** exists as both diamagnetic polymorph-

### Scheme 4. Synthesis of CAAC-Based Silylone **14**



II (major component) and paramagnetic polymorph-I with two unpaired electrons residing at the carbene carbon atoms (as shown in Scheme 4).<sup>45</sup>

**14** exhibits an obviously bent  $\text{C}_2\text{Si}$  core (Figure 4). The C–Si–C angle of  $117.44(8)^\circ$  (av) of **14** is more acute than that of





**Figure 4.** Molecular structure of **14**.

Kira's “trisilaallene” [ $136.49(6)^\circ$ ].<sup>43</sup> Although the Si–C bonds in **14** [ $1.8414(18) \text{ \AA}$ , av] are obviously longer than Si=C double bonds ( $1.702\text{--}1.775 \text{ \AA}$ ),<sup>46</sup> the  $\pi$ -type lone pair orbital at the central silicon atom (HOMO) of **14** involves notable Si–C  $\pi$  bonding. While **14** is EPR-silent, computations show that **14**, with low electronic excitation energy, has a biradicaloid character.<sup>44</sup> Subsequently, the So laboratory synthesized a mixed carbene-based analog of **14**, in which the silicon atom is embraced by both  $^{\text{Me}}\text{CAAC}$  and  $\text{NHC}^{\text{iPr}}$  [i.e.,  $\text{C}\{(\text{Pr})\text{NC}(\text{Me})\}_2$ ].<sup>47</sup> While being close to the reported Si=C double bonds ( $1.702\text{--}1.775 \text{ \AA}$ ),<sup>46</sup> the Si– $C_{\text{CAAC}}$  bond in this compound [ $1.792(4) \text{ \AA}$ ] is clearly shorter than the Si– $C_{\text{NHC}}$  bond [ $1.957(5) \text{ \AA}$ ]<sup>47</sup> and the Si– $C_{\text{CAAC}}$  bonds in **14** [ $1.8414(18) \text{ \AA}$ , av].<sup>44</sup> Both experimental and theoretical data suggest that this ( $^{\text{Me}}\text{CAAC}$ )Si( $\text{NHC}^{\text{iPr}}$ ) species can be described as a bent silaallene with a perturbed electronic structure.


Soon after compound **14** was reported, a cyclic germylone (**16**) was synthesized by Driess via low-temperature sodium naphthalenide (NaNp) reduction of the bis-NHC-complexed  $\text{GeCl}_2^+$  cation (**15**) (Scheme 5).<sup>48</sup> **15** was synthesized by reaction of the corresponding bis-NHC ligand with  $\text{GeCl}_2 \cdot \text{dioxane}$ .

X-ray structural analysis<sup>48</sup> (Figure 5) reveals that the six-membered  $\text{C}_3\text{N}_2\text{Ge}$  ring in **16** adopts a boat conformation with an acute C–Ge–C angle of  $86.6(1)^\circ$ . The Ge–C bonds in **16** [ $1.962(2)$  and  $1.967(2) \text{ \AA}$ ] are ca.  $0.10 \text{ \AA}$  shorter than those for **15** [ $2.058(3) \text{ \AA}$ , av] and ca.  $0.15 \text{ \AA}$  shorter than that in  $\text{NHC}^{\text{Dipp}}\text{-GeCl}_2$  [ $2.110(4) \text{ \AA}$ ].<sup>49</sup> While the HOMO–1 of **16** represents a  $\sigma$ -type lone pair orbital at the germanium atom, the HOMO of **16** corresponds to a  $\pi$ -type lone pair orbital of the germanium atom (with Ge–C  $\pi$  bonding interactions).<sup>48</sup>

## Scheme 5. Synthesis of Cyclic Germlyone 16

Figure 5. Molecular structure of **16**.

This  $\pi$  interaction is consistent with the relatively short Ge–C bonds in **16**. The large second proton affinities (PAs) of **14** (186.7 kcal mol<sup>-1</sup>)<sup>44</sup> and **16** (175.0 kcal mol<sup>-1</sup>)<sup>48</sup> suggest that these two molecules may be classified as divalent E(0) compounds (E = Si for **14**; Ge for **16**). Both the germanium analogue<sup>50</sup> of **14** and silicon analogue<sup>51</sup> of **16** were subsequently obtained. Notably, compound **17** (i.e., the silicon analogue of **16**)<sup>51</sup> and the recently reported mesoionic carbene-stabilized silylone (**18**)<sup>52</sup> demonstrate the capability of bonding to two metal centers (Scheme 6).<sup>22,52</sup> By

Scheme 6. Synthesis of **17**-(ZnCl<sub>2</sub>)<sub>2</sub>(THF) and **18**-(CuCl)<sub>2</sub> Binuclear Complexes (Dipp = 2,6-Diisopropylphenyl)


comparison to that of the CAAC-based silylone **14** (66.7 ppm in C<sub>6</sub>D<sub>6</sub>),<sup>44</sup> the <sup>29</sup>Si NMR resonances of **17** (-80.1 ppm in C<sub>6</sub>D<sub>6</sub>)<sup>51</sup> and **18** (-54.6 ppm, in the solid state)<sup>52</sup> are obviously higher-field shifted, indicating the presence of more electron-rich silicon cores in NHC and mesoionic carbene-based silylones. This should be ascribed to the weaker  $\pi$ -electron accepting capabilities of NHCs and mesoionic carbenes than CAACs.<sup>19,53</sup> As a result, the electron-rich silylone cores of **17** and **18** favor the formation of the corresponding binuclear complexes.

Imino-NHC<sup>54</sup> and diiminoNHC<sup>55</sup> ligands have also been utilized in accessing cyclic germlylones. While NHC-stabilized acyclic germlylones have not been reported, Jana synthesized

(L<sub>2</sub>)<sub>2</sub>Ge(0)[Fe(CO)<sub>4</sub>]<sub>2</sub> complexes (L = NHC ligands) using a “push–pull” stabilization strategy, wherein the germlylone center serves as a double  $\sigma$ -donor (coordinating to two Fe(CO)<sub>4</sub> moieties).<sup>56</sup>

Considering their electron-rich E(0) (E = Si or Ge) centers, both silylones and germlylones may be expected to demonstrate some utility in small molecule activation. To this end, reaction of **17** with CO<sub>2</sub> yielded the bis-NHC<sup>Dipp</sup>-complexed silicon decarbonate (**19**) (Scheme 7).<sup>57</sup> Although

## Scheme 7. Cyclic Silylone-Based Small Molecule Activation (Dipp = 2,6-Diisopropylphenyl)




the mechanism remains unclear, both L:SiO and L:SiO<sub>2</sub> (L = bis-NHC<sup>Dipp</sup>) have been proposed as possible intermediates in the synthesis of **19**. While bis-NHC<sup>Dipp</sup>-stabilized monomeric silicon disulfide (**20**) was obtained from reaction of **17** with elemental sulfur,<sup>58</sup> the Lewis acid–base adduct **17**-GaCl<sub>3</sub> was employed as a precursor to access bis-NHC-stabilized monomeric SiTe<sub>2</sub> (**21**), “push–pull” stabilized SiSe (**22**), and SiSe<sub>2</sub> (**23**) (Scheme 7).<sup>59</sup> A series of “push–pull” stabilized GeE (E = Se or Te), GeE<sub>2</sub> (E = S or Se) species have been prepared by reaction of the germanium analogue of **17**-GaCl<sub>3</sub> with the corresponding elemental chalcogens.<sup>60</sup> Notably, attempts to obtain CAAC-stabilized silicon oxides via aerial (or N<sub>2</sub>O) oxidation of CAAC-based Si(0) species were unsuccessful, resulting in both SiO<sub>2</sub> and N-aryl amide derivatives (CAAC=O).<sup>61</sup> In the aerial oxidation process, splitting of O<sub>2</sub> involves not only the electron-rich silicon(0)



atom but also the electrophilic CAAC carbene centers. Thus, electrophilic carbene-stabilized Si(0) species may not be an ideal platform to access the corresponding silicon oxide complexes. Despite encouraging computational efforts,<sup>62,63</sup> carbene-stabilized stannylenes and plumblylenes have not been experimentally realized. The recent synthesis of bis(silylene)- or bis(germylene)-stabilized plumblylenes may inspire synthetic chemists to finally prepare the corresponding *carbene* analogs.<sup>64,65</sup>

Dutton's computations concerning carbene-stabilization of monatomic Be(0) and diatomic Be<sub>2</sub>(0) species<sup>66</sup> suggested that carbene-complexed beryllium(0) complexes are considerably more stable than the corresponding magnesium(0) analogues. Braunschweig reported the first carbene-stabilized beryllium(0) complexes, (CAAC)<sub>2</sub>Be(0) (**26** and **27**) in 2016, via the potassium graphite reduction of CAAC-BeCl<sub>2</sub> complexes (**24** and **25**), respectively (Scheme 8).<sup>67</sup>

### Scheme 8. Synthesis of CAAC-Stabilized Be(0) Complexes (**26** and **27**) (Dipp = 2,6-Diisopropylphenyl)



The solid-state structure of **26** (Figure 6) shows that two coplanar C<sub>4</sub>N rings are bridged by a linear C<sub>carbene</sub>–Be–C<sub>carbene</sub>




Figure 6. Molecular structure of **26**.


core, which, coupled with the short Be–C<sub>carbene</sub> bonds (1.662 Å (av) for **26**; vs 1.779 Å for **24**), suggest considerable Be–C<sub>carbene</sub> π bonding interactions.<sup>67</sup> The Be–C<sub>carbene</sub> bonding in **26** and **27** is best described as a combination of donor–acceptor interactions between ground state singlet carbene ligands and the central Be(0) atom (in a 1s<sup>2</sup>2s<sup>0</sup>2p<sup>2</sup> electronic configuration). Consequently, a 3c-2e π bond is formed through the C<sub>carbene</sub>–Be–C<sub>carbene</sub> core. The strong π-accepting capabilities of CAACs are critical in stabilizing the highly reactive single Be(0) atom. However, this serves to diminish the reactivity of the beryllium(0) centers in **26** and **27**. For example, while reacting with CO<sub>2</sub> and elemental selenium, giving zwitterionic <sup>Me</sup>CAAC:CO<sub>2</sub> and selenone, respectively, **26** was shown to be inert toward dihydrogen, boranes, borohydrides, and bulky alcohols.<sup>67</sup>

### ■ CARBENE-STABILIZED E<sub>2</sub>(0) SPECIES

Seminal advances in low-oxidation state silicon chemistry over the past four decades necessarily include West's 1981 synthesis of the first disilene,<sup>68</sup> R<sub>2</sub>Si=SiR<sub>2</sub> (R = 2,4,6-(CH<sub>3</sub>)<sub>3</sub>C<sub>6</sub>H<sub>2</sub>), and Sekiguchi's 2004 report of the first disilyne,<sup>69</sup> R-Si≡Si-R (R = Dis<sub>2</sub>PrSi, Dis = bis(trimethylsilyl)methyl). The formal oxidation states of the silicon atoms in the disilene and disilyne are +2 and +1, respectively. In subsequent years a question was asked with increasing frequency: Was it possible to push the boundaries even further and prepare a compound containing a disilicon core with both silicon atoms in the formal oxidation state of zero? Relatedly, the highly reactive diatomic Si<sub>2</sub>(0) molecule has a triplet ground state (X<sup>3</sup>Σ<sub>g</sub><sup>-</sup>) and has been probed in the gas phase and in argon matrices.<sup>70</sup> The key role of elemental silicon in the semiconductor industry, coupled with its unique utility in organosilicon synthesis, only added to the allure of the elusive Si<sub>2</sub>(0) species.

Carbene-stabilization of disilicon, Si<sub>2</sub>(0), was experimentally realized by this laboratory in 2008 via the potassium graphite reduction of L:SiCl<sub>4</sub> (L = NHC<sup>Dipp</sup>) (**28**) (Scheme 9).<sup>71</sup>

### Scheme 9. Synthesis of Carbene-Stabilized Disilicon (**28** and **29**)



Schreiner subsequently suggested that London dispersion forces critically contribute to the thermodynamic stability of **28**, as well as many other bulky carbene-stabilized reactive main-group species.<sup>72</sup> One-electron oxidation of **28** by [Fe(C<sub>5</sub>Me<sub>5</sub>)<sub>2</sub>]<sup>+</sup>[B(Ar<sup>F</sup>)<sub>4</sub>]<sup>-</sup> (Ar<sup>F</sup> = C<sub>6</sub>H<sub>3</sub>-3,5-(CF<sub>3</sub>)<sub>2</sub>) results in [28]<sup>•+</sup> [B(Ar<sup>F</sup>)<sub>4</sub>]<sup>-</sup>.<sup>73</sup> NHC<sup>Dipp</sup>-stabilized Si<sup>I</sup>Cl<sub>2</sub> and Si<sup>II</sup>Cl<sub>2</sub> were also prepared by changing solvent and the reaction stoichiometry.<sup>71,74</sup> In 2014, Roesky synthesized the <sup>Cy</sup>CAAC-based disilicon (**29**) via potassium graphite reduction of <sup>Cy</sup>CAAC:SiCl<sub>4</sub> (Scheme 9).<sup>75</sup>

X-ray structural analysis<sup>71</sup> (Figure 7) shows that the Si=Si double bond distance in **28** (2.2294(11) Å) compares well to the experimental value (2.246 Å) of Si<sub>2</sub> (obtained from photoelectron spectroscopic studies).<sup>76</sup> The trans-bent geometry around the Si<sub>2</sub> core [C–Si–Si angle = 93.57(11)°], the single Si–C<sub>NHC</sub> bond [1.9271(15) Å], and the perpendicularity of the Si=Si vector to the imidazole plane are consistent with the silicon atoms in **28** residing in the formal oxidation state of zero. In contrast, if the silicon atoms in **28** reside in the +2 oxidation state, the molecule would be expected to exhibit a linear C<sub>NHC</sub>=Si=Si=C<sub>NHC</sub> core with short C=Si double




Figure 7. Molecular structure of 28.

bonds.<sup>20</sup> Computations of the simplified model 28-Ph (Figure 8) show that the HOMO and HOMO-1 correspond to the





Figure 8. HOMO, HOMO-1, and HOMO-2 molecular orbitals of the simplified 28-Ph model.

silicon-silicon  $\pi$ - and  $\sigma$ -bonds, respectively, while HOMO-2 represents one of the two lone electron pair orbitals of the  $\text{Si}_2(0)$  core.<sup>71</sup>

Due to the strong  $\pi$ -accepting capability of CAAC ligands, the lone electron pairs of silicon atoms in 29 back-donate significantly to each carbene carbon atom. Consequently, 29 exhibits shortened Si-C bonds [1.887(4) Å] and broadened C-Si-Si angles [from 101.22(13)° to 105.14(13)°] by comparison with those of 28 [ $d_{\text{Si-C}} = 1.9271(15)$  Å; C-Si-Si angle = 93.57(11)°].<sup>75</sup> Additionally, the <sup>29</sup>Si NMR resonance of 29 (249.1 ppm) is considerably shifted downfield as compared to that of 28 (224.5 ppm).

The coordinate bond essence of the Si-C<sub>NHC</sub> bonds in 28 was further confirmed by the reaction of 28 with the lithium dithiolene radical 30\*.<sup>77</sup> This reaction resulted in the release of the N-heterocyclic carbenes from the silicon atoms and subsequent formation of the dianionic silicon(IV) tris(dithiolene) complex 31 (Scheme 10).<sup>78</sup> Due to the redox-

#### Scheme 10. Carbene-Stabilized $\text{Si}_2(0)$ (28) as a Silicon-Transfer Agent




active character of the dithiolene ligand, the silicon atom is oxidized from the oxidation state of zero (in 28) to +4 (in 31). 28 acts as a silicon-transfer agent in this transformation, which joins a group of single-E-species-transfer reactions (such as E = N,<sup>79</sup> P,<sup>80,81</sup> Si,<sup>82</sup> Al,<sup>83</sup> and Pb<sup>65</sup>).

In contrast to the silicon-silicon doubly bonded disilenes, carbene-stabilized disilene(0) species contain one Si=Si

double bond in addition to two silicon-based lone pairs. The Si=Si double bond and the silicon-based lone pairs may function as  $\pi$ - and  $\sigma$ -donors, respectively. Thus, carbene-stabilized  $\text{Si}_2(0)$  species are expected to exhibit considerably different coordination behavior from disilenes.

Reaction of 28 with CuCl in toluene at -78 °C gave the 1:1 adduct (32) (Scheme 11a).<sup>84</sup> Perhaps due to steric repulsions,

#### Scheme 11. (a) Synthesis of 32 and 33, (b) $\sigma$ - $\pi$ Interconversion of 32 in Solution, (c) Topomerization of 33 in THF-*d*<sub>8</sub> Solution (L = NHC<sup>Dipp</sup>, Dipp = 2,6-Diisopropylphenyl)




attempts to isolate the 1:2 adduct [ $28:(\text{CuCl})_2$ ] repeatedly proved unsuccessful. X-ray structural analysis of 32 shows that one of the two silicon atoms is  $\sigma$ -bonded to one CuCl species. The Si-Cu bond distance in 32 [2.2081(9) Å] is marginally shorter than those in lithium bis(disilyl)cuprate [2.2412(8) and 2.2458(8) Å].<sup>85</sup> Computations revealed that the Si-Cu single bond (WBI = 0.66) in 32 is highly polarized toward silicon (78%). Notably, the Si=Si double bond distance in 32 [2.2061(12) Å] is similar to that in 28 [2.2294(11) Å],<sup>71</sup> indicating that in the solid state the silicon-silicon  $\pi$ -bond in 32 is largely not involved in the coordination to CuCl. Indeed, the simplified 32-Me model (NHC = :C[N(Me)CH]<sub>2</sub>) in  $\pi$ -bonding mode with CuCl has an elongated silicon-silicon bond (2.295 Å), which is similar to those for disilene-transition metal  $\pi$  complexes.<sup>86</sup> However, 32 only shows a singlet <sup>29</sup>Si NMR resonance in solution. In addition, the two carbenes are chemically equivalent in both <sup>1</sup>H and <sup>13</sup>C NMR spectra of 32. These NMR spectroscopic data, coupled with the results from variable-temperature (VT) <sup>1</sup>H NMR experiments, reveal that, in solution, 32 may either exist as a  $\pi$ -complex isomer or rapidly equilibrate at room temperature via a  $\pi$ -complex intermediate (Scheme 11b).<sup>84</sup> Computations show that the  $\pi$ -complex of 32-Me model (optimized in C<sub>2</sub> symmetry) is only 0.2 kcal/mol higher in energy than the corresponding  $\sigma$ -complex. The  $\sigma$ - $\pi$  interconversion of 32 (Scheme 11b) is significant since  $\sigma$ - $\pi$  rearrangements of organotransition-metal complexes are particularly important in catalytic processes.<sup>87</sup> Filippou synthesized 33 (the protonated product of 28) by

reaction of **28** with  $[\text{H}(\text{Et}_2\text{O})_2]\{\text{B}(\text{Ar}^{\text{F}})_4\}$  ( $\text{Ar}^{\text{F}} = \text{C}_6\text{H}_3\text{-}3,5\text{-}(\text{CF}_3)_2$ ) (Scheme 11a).<sup>88</sup> The dynamic NMR studies in solution revealed the topomerization (degenerate isomerization) of the  $\sigma$ -bonded tautomers of **33**, proceeding via a  $\pi$ -bonded isomer intermediate ( $\text{NHC}^{\text{Dipp}}$ -based disilahlidronium ion) (Scheme 11c). The same intramolecular topomerization phenomenon was also observed for the  $[\text{L}:(\text{I})\text{Si} = \text{Si}:\text{L}]^+$  cation ( $\text{L} = \text{NHC}^{\text{Dipp}}$ ).<sup>89</sup>

Carbene-stabilized disilicon(0), **28**, may serve as an effective platform to access unusual silylene complexes. For example, the 1:4 reaction of **28** with  $\text{BH}_3 \cdot \text{THF}$  in toluene resulted in the cleavage of the  $\text{Si}=\text{Si}$  double bond, giving **34** in 72% yield (Scheme 12).<sup>90</sup> The highly reactive  $\text{SiH}_2$  species has been


### Scheme 12. Borane-Mediated Cleavage of **28**



observed as an intermediate in the chemical vapor deposition of silicon film via  $\text{SiH}_4$  pyrolysis.<sup>91</sup> **34** represents the first “push–pull”-stabilized parent silylene ( $\text{SiH}_2$ ), wherein the  $\text{SiH}_2$  unit accepts one pair of electrons from the NHC ligand, while donating one electron pair to the  $\text{NHC}:\text{Si}(\text{H})(\text{B}_3\text{H}_7)\text{BH}_2$  moiety. In the presence of a small amount of free NHC ligand (**28** to NHC = 5:1), the parallel reaction gives a mixture containing both **34** (30% yield) and **35** (28% yield) (Scheme 12).<sup>90</sup> Compound **35** is a “push–pull”-stabilized three-membered cyclosilylene, wherein the silylene center accepts an electron pair from the carbene ligand, while donating a pair of electrons to the  $\text{BH}_3$  unit. The formation of **35** may involve cycloaddition of the silicon(0) atom of the  $\text{NHC}:\text{Si}(\text{BH}_3)$  intermediate to the  $\text{C}=\text{C}$  backbone of the NHC ligand.

Room temperature reaction of **28** with iron pentacarbonyl (in a 1:1 ratio) in toluene gives the tetracarbonyliron adduct, **36**, as a dark purple solid in 81% yield (Scheme 13).<sup>92</sup> X-ray structural analysis of **36** shows one silicon atom is  $\sigma$ -bonded to one  $\text{Fe}(\text{CO})_4$  moiety. In contrast to **32**,<sup>84</sup> which only shows one singlet  $^{29}\text{Si}$  NMR resonance (226.7 ppm in  $\text{C}_6\text{D}_6$ ), **36** exhibits two  $^{29}\text{Si}$  NMR resonances (142.5 and 201.3 ppm in  $\text{THF-d}_8$ ), indicating the presence of the asymmetrical structure of **36** not only in the solid state but also in solution. The lack of dynamic complexation behavior of **36** in solution may be due to the steric bulk of the  $\text{Fe}(\text{CO})_4$  fragment. Compound **37** can be prepared either through the 1:1 reaction of **36** with  $\text{Fe}(\text{CO})_5$  at 100 °C or by direct reaction of **28** with excess  $\text{Fe}(\text{CO})_5$  (Scheme 13).<sup>92</sup> The **28**-to-**37** conversion involves the insertions of both CO and  $\text{Fe}_2(\text{CO})_6$  into the two  $\text{NHC}^{\text{Dipp}}$ -stabilized silicon atoms. Furthermore, reaction of **36** with pyridine hydrochloride ( $\text{HCl} \cdot \text{NC}_5\text{H}_5$ ) gave a “push–pull”-stabilized parent monochlorosilylene  $[\text{Si}(\text{H})\text{Cl}]$  (**38**)

### Scheme 13. Synthesis of **36–38** (Dipp = 2,6-Diisopropylphenyl)




(Scheme 13), in which two  $\text{NHC}^{\text{Dipp}}$ -complexed  $:\text{Si}(\text{H})\text{Cl}$  silylene units donate electron pairs to an  $\text{Fe}(\text{CO})_3$  moiety.<sup>93</sup>

Unlike naturally occurring stable silica ( $\text{SiO}_2$ ), which consists of a covalent  $\text{Si}-\text{O}-\text{Si}$  network, molecular  $\text{SiO}_2$  and other small silicon oxides are highly reactive. Molecular  $\text{SiO}_2$  has been detected in solid noble-gas matrices<sup>94,95</sup> and gas-phase reactions<sup>96</sup> and explored by photoelectron spectroscopy.<sup>97</sup> By allowing **28** to react with  $\text{O}_2$  and  $\text{N}_2\text{O}$ , this laboratory obtained two novel molecules— $\text{NHC}^{\text{Dipp}}$ -stabilized  $\text{Si}_2\text{O}_4$  (**39**) and  $\text{Si}_2\text{O}_3$  (**40**), respectively (Scheme 14).<sup>98</sup> X-ray structural analysis shows that the  $\text{Si}_2\text{O}_4$  core of **39** adopts  $\text{C}_{2h}$  symmetry due to carbene coordination. While each silicon(IV) atom in **39** is capped by a terminal oxygen atom with a 1.5260(14) Å  $\text{Si}-\text{O}_{\text{terminal}}$  bond (bearing modest double-bond character), these two silicon(IV) atoms are also bridged by two additional oxygen atoms with  $\text{Si}-\text{O}_{\text{bridge}}$  single bonds (1.675 Å, av). The  $\text{Si}-\text{C}_{\text{NHC}}$  bond distance in **39** [1.9259(17) Å] compares well to that in **28** [1.9271(15) Å].<sup>71</sup> Regarding the  $\text{Si}_2\text{O}_3$  core of **40**, the two silicon(III) atoms are bridged by one oxygen atom while retaining a silicon–silicon single bond [2.2405(14) Å].<sup>98</sup> Further oxidation of **40** by  $\text{CO}_2$  gives a carbene-stabilized silicon–carbon mixed oxide ( $\text{SiO}_2$ )<sub>2</sub> $\text{CO}_2$  (**41**), which can also be directly prepared via  $\text{CO}_2$  oxidation of **28** (Scheme 14).<sup>99</sup> By allowing **28** to react with elemental tellurium, So and co-workers synthesized not only the tellurium analogues of **39** and **40** but also  $\text{NHC}^{\text{Dipp}}$ -stabilized  $\text{Si}_2\text{Te}_2$  and  $\text{Si}_2\text{Te}_2\text{S}$  clusters.<sup>100</sup> Notably, the  $\text{NHC}$ -stabilized  $\text{Si}_2\text{Te}_2$  may exist in two isomeric forms [i.e.,  $\text{NHC}(\text{Te})\text{Si}=\text{Si}(\text{Te})\text{NHC}$  and  $\text{NHC}(\text{Te})\text{Si}(\mu\text{-Te})\text{SiNHC}$ ]. The stability sequence of these  $\text{NHC}$ -complexed  $\text{Si}_2\text{Te}_n$  ( $n = 2, 3$ , and 4) clusters are  $\text{Si}_2\text{Te}_4 > \text{Si}_2\text{Te}_3 > (\text{Te})\text{Si}(\mu\text{-Te})\text{Si} > (\text{Te})\text{Si}=\text{Si}(\text{Te})$ .<sup>100</sup>

By utilizing CAAC-stabilized disilicon complexes (**29** and **42**)<sup>75,101</sup> (Scheme 14), Roesky synthesized CAAC-stabilized  $\text{Si}_2\text{S}_4$  (**43**)<sup>102</sup> and  $\text{Si}_2\text{Se}_4$  (**44** and **45**).<sup>101</sup> In contrast to **43–45**, which are stable for months under an inert atmosphere at room temperature, crystals of  $\text{NHC}^{\text{Dipp}}$ -stabilized  $\text{Si}_2\text{O}_4$  (**39**)


**Scheme 14. Carbene-Stabilized-Disilicon-Mediated Small Molecule Activation (Dipp = 2,6-Diisopropylphenyl, Cy = Cyclohexyl)**



and  $\text{Si}_2\text{O}_3$  (40) slowly decompose, even under argon protection. Storage of 39 and 40 at low temperature ( $-40^\circ\text{C}$ ) would obviously increase the stability of these two compounds. The high reactivity of 39 and 40 indicates that these compounds may be employed to transfer and integrate the “ $\text{Si}_2\text{O}_3$ ” or “ $\text{Si}_2\text{O}_4$ ” units into various inorganic and organometallic substrates. CAAC-stabilized silicon oxides, via oxidation of 29 and 42, have not been reported. When exposed to air, 42 decomposed to  $^{\text{Me}}\text{CAAC}=\text{O}$  and  $\text{SiO}_2$ .<sup>61,101</sup>

Employing  $\text{RMg-MgR}$  ( $\text{R} = \beta$ -diketiminate ligand) as a two-center, two-electron reducing agent to react with  $\text{NHC}^{\text{Dipp}}\text{ECl}_2$  [ $\text{E} = \text{Ge}$  (46),  $\text{Sn}$  (47)], Jones and Stasch synthesized carbene-stabilized  $\text{Ge}_2(0)$  (48) and  $\text{Sn}_2(0)$  (49), respectively (Scheme 15).<sup>49,103</sup> Similar to carbene-stabilized  $\text{Si}_2(0)$  (28),<sup>71</sup> both 48 and 49 adopt the trans-bent geometry around the  $\text{E}=\text{E}$  double bond [ $d_{\text{E}=\text{E}} = 2.3490(8) \text{ \AA}$ ,  $\text{E} = \text{Ge}$  (48);  $d_{\text{E}=\text{E}} = 2.7225(5) \text{ \AA}$ ,  $\text{Sn}$  (49)]. The carbene  $^{13}\text{C}$  NMR resonances of  $\text{NHC}^{\text{Dipp}}$ -stabilized  $\text{Si}_2(0)$  (28) (196.3 ppm),  $\text{Ge}_2(0)$  (48) (203.3 ppm), and  $\text{Sn}_2(0)$  (49) (210.3 ppm) are shifted downfield upon descending the group, thus indicating a weaker  $\text{E}-\text{C}_{\text{NHC}}$  bonding interaction with increasing atomic mass of E.

**Scheme 15. Synthesis of NHC-Stabilized  $\text{Ge}_2(0)$  (48),  $\text{Sn}_2(0)$  (49), and “Push–Pull”-Stabilized  $\text{Ge}_2(0)$  (51)**



Donor–acceptor-stabilized  $\text{Ge}_2(0)$  species (51) was synthesized by Scheschkewitz via potassium-graphite reduction of  $\text{NHC:Ge(Cl)}_2[\text{Fe(CO)}_4]$  ( $\text{NHC} = :\text{C}\{(\text{Pr})\text{NC}(\text{Me})\}_2$ ) in THF (Scheme 15).<sup>104</sup> X-ray structural analysis of 51 shows that while one  $\text{Fe(CO)}_4$  moiety is terminally bonded to a germanium atom with a  $\text{Ge-Fe}$   $\sigma$ -bond [ $2.4112(3) \text{ \AA}$ ], the other  $\text{Fe(CO)}_4$  unit acts as a bridge between the two germanium atoms [bridging  $\text{Ge-Fe}$  bond distances =  $2.6292(3) \text{ \AA}$ , av]. Consequently, the  $\text{Ge-Ge}$  bond in 51 [ $2.4442(2) \text{ \AA}$ ] is slightly longer (ca.  $0.1 \text{ \AA}$ ) than that in  $\text{NHC}^{\text{Dipp}}$ -stabilized  $\text{Ge}_2(0)$  (48) [ $2.3490(8) \text{ \AA}$ ].<sup>49</sup> Computations support that 51 has a metallacyclopropane-type bonding motif, according to the Dewar–Chatt–Duncanson model. Both the NMR spectral data and theoretical calculations of 51 imply a possible degenerate equilibrium between 51 and 51' through an isomeric species with two terminal  $\text{Fe(CO)}_4$  units




[50] (Scheme 15).<sup>104</sup> Indeed, [50] is only slightly higher in free energy (by only 7.4 kcal mol<sup>-1</sup>) than 51. The interaction between the digermanium unit and the bridging Fe(CO)<sub>4</sub> fragment is weak and should thus involve certain  $\pi$ -complex character. Room temperature reaction of 51 with propylene sulfide (in toluene) gives a digermathirane (52) containing a Ge<sub>2</sub>S three-membered ring (Scheme 15).<sup>104</sup>

In contrast to the NHC-stabilized heavier E<sub>2</sub>(0) (E = Si, Ge, and Sn) congeners, wherein the E<sub>2</sub>(0) cores exhibit trans-bent geometries [C<sub>NHC</sub>-E-E angle = 93.57(11)°, E = Si (28);<sup>71</sup> 89.87(8)°, E = Ge (48);<sup>49</sup> 91.82(8)°, E = Sn (49)],<sup>103</sup> the NHC-based dicarbon derivative was predicted in 2012 by Dutton and Wilson to exist as a linear cumulene.<sup>105</sup> An experimental attempt to obtain the NHC-based dicarbon, through deprotonation of a doubly protonated precursor 53, proved unsuccessful, affording only the reduced product 54 (Scheme 16).<sup>106</sup> CAAC-based cumulenes 55 and 56 were independently synthesized by the laboratories of Roesky and Bertrand, respectively (Scheme 16).<sup>107,108</sup> In addition, Kinjo reported a 4-pyridylidene-based cumulene (57) (Scheme 16).<sup>109</sup> The bonding analysis, using charge and energy decomposition methods, revealed that (CAAC<sup>Me</sup>)<sub>2</sub>C<sub>2</sub> and (DAC<sup>Me</sup>)<sub>2</sub>C<sub>2</sub> (DAC = diamidocarbene) possess genuine cumulene C<sub>4</sub> cores due to the electron-sharing bonding between quintet L<sub>2</sub> (L = CAAC<sup>Me</sup> and DAC<sup>Me</sup>) and quintet C<sub>2</sub> fragments.<sup>110</sup> However, the bonding in (NHC<sup>Me</sup>)<sub>2</sub>C<sub>2</sub> and (SNHC<sup>Me</sup>)<sub>2</sub>C<sub>2</sub> (SNHC = saturated NHC) appears to have been based on a combination of dative and electron-sharing interactions between doublet L<sub>2</sub><sup>+</sup> (L = NHC<sup>Me</sup> and SNHC<sup>Me</sup>) and doublet C<sub>2</sub><sup>-</sup> moieties.<sup>110</sup> In contrast to the linear “cumulene” cores of 55–57, a monoligated L:C<sub>2</sub> complex (58) [L: = (NHC<sup>Dipp</sup>=N)<sub>2</sub>(Me)P] was recently synthesized by Ong, using a sterically demanding phosphine ligand (Scheme 16).<sup>111</sup> Notably, these phosphine-stabilized dicarbon complexes have been utilized as supporting ligands in transition-metal catalysis.<sup>112</sup>


Following our discovery of carbene-stabilized disilicon (28), we extended the carbene-stabilization strategy to the group 15 elements. The potassium graphite reduction of NHC<sup>R</sup>:ECl<sub>3</sub> in THF gives the corresponding carbene-stabilized E<sub>2</sub>(0) species (E = P, R = Dipp, 59; E = P, R = Mes, 60; E = As, R = Dipp, 61) (Scheme 17).<sup>113,114</sup> Alternatively, 59 may also be accessed by using Na(OCP) as a phosphorus-transfer agent.<sup>115</sup>

The dominant structural feature of 59 and 60 is the phosphorus–phosphorus single bond [d<sub>P–P</sub> = 2.2052(10) Å, 59; d<sub>P–P</sub> = 2.1897(11) Å, 60] (Figure 9).<sup>113</sup> As a result, the steric demands of the carbenes may significantly affect the conformations of these complexes. While the P<sub>2</sub> core in 59 exhibits a trans-bent geometry with the C–P–P–C torsion angle of 180.0°, the corresponding P<sub>2</sub> core in 60 adopts a gauche conformation (the C–P–P–C torsion angle = 134.1°). The P–C<sub>NHC</sub> bond distances of 59 [1.7504(17) Å] and 60 [1.754(3) Å] are between P=C double bond distances of the nonconjugated phosphalkenes<sup>116</sup> (1.65–1.67 Å) and typical P–C single bond distances (such as that of NHC<sup>Dipp</sup>:PCl<sub>3</sub> [1.871(11) Å]).<sup>117</sup> The WBI of 59 (1.40) suggests modest double bond character of the P–C<sub>NHC</sub> (due to the back-donation of the electron pair of the phosphorus atom to the p orbital of the carbene carbon atom). However, the high-field <sup>31</sup>P NMR resonances for 59 (–52.4 ppm) and 60 (–73.6 ppm) support the presence of electron-rich bis-(phosphinidene) cores in these two complexes. Isostructural to 59, carbene-stabilized diarsenic, 61, containing a singly


### Scheme 16. Synthesis of Cumulenes (55–57) and Monoligated C<sub>2</sub> Complex (58) (Dipp = 2,6-Diisopropylphenyl)



### Scheme 17. Synthesis of NHC-Stabilized P<sub>2</sub>(0) (59 and 60) and As<sub>2</sub>(0) (61)




bonded As<sub>2</sub>(0) core [d<sub>As–As</sub> = 2.442(1) Å], may be described as a carbene-complexed diarsinidene.<sup>114</sup>

Figure 9. Molecular structures of **59** and **60**.

Reaction of **59** with excess  $\text{BH}_3 \cdot \text{THF}$  gave the boronium complex **62** in 85% yield (Scheme 18), wherein the carbene-

Scheme 18. Synthesis of **62**–**66** (L = NHC<sup>Dipp</sup>; Dipp = 2,6-Diisopropylphenyl)




stabilized  $\text{P}_2(0)$  core serves as a bidentate ligand to bind to a  $\text{BH}_2^+$  cation.<sup>118</sup> The  $^1\text{H}$  NMR spectroscopy suggests an equilibrium between **59** and **62**, which favors the formation of **62**. Addition of excess of  $\text{BH}_3 \cdot \text{THF}$  was shown to significantly diminish the dissociation of **62**. While reduction of **59** with lithium metal in THF gave the  $\text{C}_4$ -lithiated NHC-stabilized parent phosphinidene (PH) (**63**),<sup>117</sup> oxidation of **59** with dioxygen in toluene results in NHC-stabilized  $\text{P}_2\text{O}_4$  (**64**) (Scheme 18).<sup>119</sup> In contrast to the highly reactive carbene-stabilized  $\text{Si}_2\text{O}_4$  (**39**) and  $\text{Si}_2\text{O}_3$  (**40**),<sup>98</sup> complex **64** is air-stable. The splitting of triplet  $\text{O}_2$  by the singlet  $\text{P}_2(0)$  core of **59** may have involved single-electron transfer processes.<sup>120</sup> The  $\text{P}_2\text{O}_4$  core in **64** exists as a  $\text{PO}_2$  dimer containing a P–P single bond [2.310(2) Å]. Notably, free  $\text{P}_2\text{O}_4$  energetically favors an oxo-bridged and nonplanar  $\text{O}_2\text{POPO}$  isomer<sup>121,122</sup> (with  $\text{C}_s$  symmetry) rather than the symmetric  $\text{O}_2\text{P}=\text{PO}_2$

dimer observed in **64**. Bertrand reported that while  $[\text{Ph}_3\text{C}]^+[\text{B}(\text{C}_6\text{F}_5)_4]^-$ -mediated one-electron oxidation of **59** provided the NHC-stabilized  $\text{P}_2^{\bullet+}$  radical cation (**65**), ferrocenium triflate mediated two-electron oxidation of **59** gave the NHC-stabilized  $\text{P}_2^{2+}$  dication (**66**) (Scheme 18).<sup>123</sup>


This laboratory subsequently discovered that the Lewis acid  $\text{GaCl}_3$  may be employed as an oxidant to convert NHC-stabilized  $\text{As}_2(0)$  (**61**) into either the corresponding monocationic diarsenic radical (**67**) or the dicationic diarsene (**68**) (Scheme 19).<sup>124</sup>

Scheme 19. Synthesis of **67** and **68** (L = NHC<sup>Dipp</sup>; Dipp = 2,6-Diisopropylphenyl)



Bertrand reported that  $\text{C}_y\text{CAAC}$  (**69**) can mediate fragmentation of white phosphorus ( $\text{P}_4$ ), giving both  $(\text{L:P})_3\text{P}$  (**70**) and  $\text{L:P}=\text{P:L}$  (**71**) (L = **69**) (Scheme 20).<sup>125</sup> The P–C bond distance of **71** [1.719(7) Å] is ca. 0.03 Å shorter than that (ca. 1.75 Å) of NHC-stabilized  $\text{P}_2(0)$  (**59** and **60**).<sup>113</sup> In


Scheme 20. Synthesis of CAAC-Based  $\text{P}_2(0)$  (**71**),  $\text{As}_2(0)$  (**73**), and  $\text{Sb}_2(0)$  (**74**) (Dipp = 2,6-Diisopropylphenyl)



contrast to the high-field  $^{31}\text{P}$  NMR resonances of **59** ( $-52.4$  ppm) and **60** ( $-73.6$  ppm), compound **71** shows a low-field  $^{31}\text{P}$  NMR resonance of  $54.2$  ppm. These structural and spectroscopic data strongly support the 2,3-diphosphabutadiene essence of **71**. While **59** can be converted to **65** and **66** via one-electron and two-electron oxidation, respectively (Scheme 18), CAAC-stabilized diphosphorus (**71**) can only be converted to the analogue of **65** via  $[\text{Ph}_3\text{C}]^+[\text{B}(\text{C}_6\text{F}_5)_4]^-$ -mediated one-electron oxidation.<sup>123</sup> The  $\text{Et}^i\text{CAAC}$ -stabilized diarsenic (**73**) was synthesized by Hudnall via potassium graphite reduction of the  $[(\text{Et}^i\text{CAAC})_2\text{As}]^+\text{Cl}^-$  precursor (**72**) (Scheme 20).<sup>126</sup> The As–C bond distance of **73** [ $1.837(5)$  Å] is ca.  $0.04$  Å shorter than that of **61** [ $1.881(2)$  Å].<sup>114</sup> The E=C (E = P, As) double bond character (in **71** and **73**) should be attributed to the increased  $\pi$ -acidic character of the CAAC ligand compared to the NHC. By allowing one equivalent of  $\text{C}^y\text{CAAC}:\text{SbCl}_3$  to react with 3 equiv of potassium graphite, Bertrand obtained  $\text{C}^y\text{CAAC}$ -stabilized  $\text{Sb}_2(0)$  (**74**) (Scheme 20).<sup>127</sup> Compound **74** contains a singly bonded  $\text{Sb}_2(0)$  core [ $d_{\text{Sb-Sb}} = 2.8125(10)$  Å], which adopts an anticlinal twisted-bent geometry [C–Sb–Sb–C torsion angle =  $122.6(4)^\circ$ ]. The Sb–C bond distances in **74** [ $2.084(11)$ – $2.088(10)$  Å] are longer than the theoretical value for the parent stiba-alkene ( $2.01$  Å).<sup>128</sup> The WBI value of the Sb–C bond in **74** ( $1.23$ ) is less than that of the As–C bond in **61** ( $1.34$ ) and that of the P–C bond in **59** ( $1.40$ ), indicating the decreased multiple bond character of the E–C<sub>carbene</sub> bonds, descending group 15. Notably, among carbene-stabilized main group  $\text{E}_2(0)$  species, **74** contains the heaviest diatomic allotrope core reported.


This laboratory has long been fascinated by the multiple bond chemistry of the group 13 elements.<sup>129,130</sup> Although our initial goal was to prepare a molecule containing a boron–boron triple bond, the potassium graphite reduction of  $\text{NHC}^{\text{Dipp}}:\text{BBr}_3$  afforded a  $\text{NHC}^{\text{Dipp}}$ -stabilized neutral diborene (**75**) (orange-red crystals) and a diborane (**76**) (colorless crystals) (Scheme 21).<sup>38</sup> The stoichiometric

### Scheme 21. Synthesis of NHC-Stabilized Neutral Diborene (75) and Diborane (76)



ratio of  $\text{NHC}^{\text{Dipp}}:\text{BBr}_3$  to  $\text{KC}_8$  has been observed to affect the yield of **75**. A higher yield (12%) of **75** was obtained with a 1:5.4 molar ratio of  $\text{NHC}^{\text{Dipp}}:\text{BBr}_3$  to  $\text{KC}_8$ .


In contrast to the four-coordinate tetrahedral boron atoms of **76** (involving a boron–boron single bond of  $1.828(4)$  Å), X-ray data<sup>38</sup> (Figure 10a) revealed the most salient structural feature of **75**<sup>38</sup>—the three-coordinate trigonal planar boron atoms constituting a boron–boron double bond—the first neutral diborene. The boron–boron double bond distance in **75** [ $1.560(18)$  Å, av] is shorter than those distances reported for diboron dianions  $[\text{Mes}_2\text{BB}(\text{Mes})\text{Ph}]^{2-}$  [ $1.636(11)$  Å]<sup>131</sup> and  $[\{\text{Ph}(\text{Me}_2\text{N})\text{BB}(\text{NMe}_2)\text{Ph}\}]^{2-}$  ( $1.627$  Å, av).<sup>132</sup> While the HOMO of the simplified **75-H** model corresponds to a B–B  $\pi$ -bonding orbital, the HOMO–1 involves mixed B–B and B–H  $\sigma$ -bonding character (Figure 10b).<sup>38</sup>



**Figure 10.** Molecular structure of **75** (a); the HOMO and HOMO–1 orbitals of the simplified **75-H** model (b).

The presence of hydrides (B–H) in **75** and **76** was presumed to be due to hydrogen abstraction from ethereal solvent in the presence of alkali metals.<sup>133</sup> Decreasing the steric bulk of the carbene ligand may aid the boron–boron bond formation and thus diminish the chance of hydrogen abstraction from the reaction media. Subsequently, this laboratory conducted potassium graphite reduction of less bulky  $\text{NHC}^{\text{Mes}}$ -complexed  $\text{BBr}_3$ . However, this reaction afforded three polymorphs of the neutral diborane  $\text{L}:\text{B}(\text{H})=\text{B}(\text{H})\text{:L}$  (L =  $\text{NHC}^{\text{Mes}}$ ) (**77**) with planar, twisted, and trans-bent structures.<sup>134</sup> In 2012, the Braunschweig laboratory made a remarkable breakthrough in this field. Following the synthesis of **78** by reacting tetrabromodiborene(4) with 2 equiv of  $\text{NHC}^{\text{Dipp}}$ ,<sup>135</sup> they conducted sodium naphthalene mediated reduction of **78** in THF at  $-78$  °C. This resulted in the  $\text{NHC}^{\text{Dipp}}$ -stabilized  $\text{B}_2(0)$  (**79**), containing the long-sought boron–boron triple bond (Scheme 22).<sup>135</sup>

### Scheme 22. Synthesis of Carbene-Based Diboryne (79) and Diboracumulene (81) (Dipp = 2,6-Diisopropylphenyl)



In the solid state (Figure 11),<sup>135</sup> **79** contains an essentially linear C–B $\equiv$ B–C core (the C–B–B–C torsion angle =  $-161.83^\circ$ ). Each boron atom is two-coordinate with a C–B–B angle of ca.  $173.0^\circ$ . The boron–boron triple bond distance of  $1.449(3)$  Å matches well with the experimental values ( $1.453$  to  $1.468$  Å) of  $\text{OCBBCO}$ .<sup>136</sup> It is interesting to compare the structural and spectroscopic data of **79** with the  $\text{NHC}^{\text{Dipp}}$ -complexed diborene (**75**) and diborane (**76**).<sup>38</sup> The B $\equiv$ B triple bond in **79** [ $1.449(3)$  Å] is ca.  $0.11$  Å shorter than the B=B double bond in **75** [ $1.560(18)$  Å, av] and ca.  $0.38$  Å shorter than the B–B single bond in **76** [ $1.828(4)$  Å]. The B–C<sub>NHC</sub> bond distances shorten in sequence: **76** [ $1.577(2)$  Å] > **75** [ $1.538(15)$  Å, av] > **79** [ $1.491(3)$  Å, av], indicating




Figure 11. Molecular structure of **79**.


increased delocalization of the  $\pi$ -electrons of boron over the B–C<sub>NHC</sub> fragments. With the decrease of the coordination number of boron, the <sup>11</sup>B NMR resonances are shifted downfield in sequence: **76** (–31.6 ppm), **75** (25.3 ppm), **79** (39.0 ppm). The Raman<sup>137</sup> and NMR<sup>138</sup> spectroscopic studies provide further evidence for the *triple bond* character of the B<sub>2</sub>(0) core in **79**.

Braunschweig also prepared the CAAC-complexed diboron **81** via sodium naphthalenide reduction of **80**, the CAAC-based analogue of **78** (Scheme 22).<sup>139</sup> The electronic properties of the carbene ligands have a remarkable effect on the bonding pattern of the C<sub>2</sub>B<sub>2</sub> cores in these carbene-stabilized B<sub>2</sub> complexes. CAACs have been reported to be stronger  $\sigma$ -donors and  $\pi$ -acceptors than NHCs.<sup>7</sup> Hence, the boron–boron bond in **81** [1.489(2) Å] is somewhat longer than that in **79** [1.449(3) Å],<sup>135</sup> yet still shorter than the reported B=B double bonds (1.56–1.71 Å).<sup>140,141</sup> Concomitantly, the B–C bonds in **81** [1.458(2) Å, av] are shorter than those in **79** [1.491(3) Å, av]. Each boron atom in **79** bears an NPA charge of –0.13, indicating the electron-rich of the B<sub>2</sub>(0) core in **79**. However, each boron atom in **81** has an NPA charge of +0.08. The 80.0 ppm <sup>11</sup>B NMR resonance of **81** is shifted significantly downfield compared to that of **79** (39.0 ppm). Thus, it may be appropriate to describe **81** as a diboracumulene. The  $\pi$ -acidity of the saturated NHC (i.e., SNHC) ligands is stronger than that of unsaturated NHCs but weaker than that of CAACs. As a result, the C<sub>2</sub>B<sub>2</sub> cores in SNHC<sup>Dipp</sup> or SNHC<sup>Dip</sup>-stabilized diborons (**82**<sup>142</sup> and **83**<sup>143</sup>) exhibit an intermediate bonding pattern in-between that of **79** and that of **81**.

Considering the well developed chemistry of transition metal alkyne complexes, particularly their pivotal role in catalytic reactions, it is logical to investigate the coordination chemistry of the lightest triple bond (i.e., the B≡B bond) in **79**. The first transition metal  $\pi$ -complexes involving boron–boron triple bonds were synthesized by Braunschweig via reactions of **79** with copper(I) species (Scheme 23).<sup>144</sup> Reaction of **79** with 2 equiv of [CuCC(SiMe<sub>3</sub>)] gave **84**, wherein the B<sub>2</sub>(0) core of **79** is  $\pi$ -bonded to two [CuCC(SiMe<sub>3</sub>)] units. Further reaction of **84** with 1 equiv (at 0 °C) or 3 equiv (at room temperature) of [CuCl(SMe<sub>2</sub>)] produced diboryne- $\pi$ -complexed [CuCl]<sub>2</sub> (**85**) or [CuCl]<sub>3</sub> (**86**), respectively. The trinuclear complex **86** was also directly prepared by reacting **79** with 4 equiv of [CuCl(SMe<sub>2</sub>)] (Scheme 23).

As a result of di- and trimetalations, both B≡B and B–C bond distances increase in sequence: **79** [ $d_{B-B}$  = 1.449(3) Å;  $d_{B-C}$  = 1.491(3) Å, av] < **84** [ $d_{B-B}$  = 1.478(2) Å;  $d_{B-C}$  = 1.534(2) Å, av] and **85** [ $d_{B-B}$  = 1.486(5) Å;  $d_{B-C}$  = 1.546(4) Å, av] < **86** [ $d_{B-B}$  = 1.526(4) Å;  $d_{B-C}$  = 1.562(3) Å]. In addition, these diboryne- $\pi$ -complexed Cu<sub>*n*</sub> (*n* = 2, 3) species display intense (for **86**) or weak (for **84** and **85**)


Scheme 23. Synthesis of **84–86** (L = NHC<sup>Dipp</sup>; Dipp = 2,6-Diisopropylphenyl)



phosphorescence in the red to near-IR region from their triplet excited states with quantum yields of up to 58%. DFT computations show that complexes **84–86** exhibit enhanced metal d orbital contributions to HOMO and HOMO–1. This leads to S<sub>1</sub> and T<sub>1</sub> with remarkable MLCT character and enables strong spin–orbit coupling for highly efficient intersystem-crossing S<sub>1</sub> → T<sub>n</sub> and phosphorescence T<sub>1</sub> → S<sub>0</sub> transitions.<sup>144</sup>

Diboryne **79** also formed  $\pi$ -complexes with main group cations. When **79** was combined with diaryltellurides, diboryne- $\pi$ -complexed RTe<sup>+</sup> cations (**87** and **88**) were obtained via nucleophilic attack by **79** on one of the two tellurium atoms of diaryltellurides (Scheme 24).<sup>145,146</sup> The B–

Scheme 24. Synthesis of **87–90** (L = NHC<sup>Dipp</sup>; Dipp = 2,6-Diisopropylphenyl)



B bonds of **87** [1.490(6) Å] and **88** [1.494(10) Å] are somewhat longer than that of **79** [1.449(3) Å], but still obviously shorter than those for diborene complexes [such as  $d_{B=B}$  = 1.560(18) Å (av) for **75**].<sup>38</sup> The elongation of the bound boron–boron bonds in **87** and **88** is limited. For example, the boron–boron bond in **87** is merely 2.8% longer than the free B≡B bond in **79**. The B–B–C angles in **87** and **88** (ca. 163–165°) are slightly bent from linearity. These structural features support assignment of **87** and **88** as  $\pi$ -complexes.<sup>146</sup> When **79** was combined with M[B(3,5-




$C_6H_3Cl_2)_4]$  ( $M = Li, Na$ ), complexes **89** and **90** were formed, respectively (Scheme 24).<sup>147</sup> In these complexes, the lithium or sodium cations are  $\pi$ -encapsulated by the diboryne unit, a rare feature for neutral molecules. In contrast to the cases of **84–88**, the B–B bond distances in **89** [1.452(6) Å] and **90** [1.459(4) Å] compare well to that of **79** [1.449(3) Å], indicating little covalent interaction between alkali metal cations and the B $\equiv$ B triple bond. Notably, encapsulation of alkali metal cations by neutral NHC-stabilized Si<sub>2</sub>(0) (**28**) has also been observed.<sup>88</sup>

With a series of carbene-stabilized diboron complexes available, Braunschweig further investigated their capacity to activate small molecules. Room temperature reaction of **79** with excess elemental sulfur and selenium resulted in isostructural **91** and **92**, respectively, via reductive insertion of elemental chalcogens into the boron–boron triple bond (Scheme 25).<sup>148</sup> Both **91** and **92** contain a [2.2.1]-bicyclic system, wherein the two boron atoms are bridged by five chalcogen atoms. These reactions involved a six-electron reduction, supporting the B $\equiv$ B triple bond character of **79**. While reactions of NHC-based diborynes with CO<sub>2</sub> provided intractable mixtures, the CO<sub>2</sub>-mediated oxidation of diboracumulene (**81**), involving an initial [2 + 2] cycloaddition, resulted in **93** (Scheme 25).<sup>149</sup> The X-ray structural analysis of **93** demonstrated that **81** can fix two CO<sub>2</sub> molecules. One CO<sub>2</sub> molecule is cleaved to form a boron-bound terminal CO. Both the released oxygen atom and the other CO<sub>2</sub> molecule insert into a B–C<sub>CAAC</sub> bond, giving a spiro five-membered C<sub>2</sub>O<sub>2</sub>B ring. In the contrast, the Si–C<sub>NHC</sub> bonds remain intact in the CO<sub>2</sub>-mediated oxidation of NHC-stabilized Si<sub>2</sub>(0) (**28**).<sup>99</sup> The low-temperature reaction of **79** with CO (in a 1:2 molar ratio) resulted in the isolation of a stable intermediate **94** [containing an unsymmetrically bridging CO between two doubly bonded boron atoms ( $d_{B=B} = 1.549(3)$  Å)]. **94** was then quantitatively converted to the NHC<sup>Dipp</sup>-stabilized bis(boralactone) (**95**) by reaction with excess CO at room temperature (Scheme 25).<sup>150</sup> However, the reaction of **81** with CO gave a stable bis(boraketene) (**96**) (Scheme 25),<sup>142</sup> which could not be converted to the CAAC-based bis(boralactone) even at 150 °C and under 50 bar of CO. When the SNHC<sup>Dipp</sup>-stabilized diboron (**82**) was combined with CO, the corresponding carbene-stabilized bis(boraketene) (**97**) and bis(boralactone) (**98**) were obtained.<sup>142</sup> While **79**, **82**, and NHC<sup>Dep</sup>-based B<sub>2</sub>(0) (**99**)<sup>137</sup> cannot undergo hydrogenation, both **83** and **81** may react with H<sub>2</sub> at 1 atm, giving SNHC<sup>Dep</sup>- and CAAC-stabilized parent diborenes (**100** and **101**), respectively (Scheme 25).<sup>143</sup> These results suggest that the electronic and steric properties of the associated carbene ligands play a critical role in carbene-diboron-mediated activation of small molecules. A recent study indicated that the reactivity of carbene-stabilized diboron complexes may be enhanced by desymmetrization (i.e., formation of a zerovalent sp–sp<sup>2</sup> diboron complex via NHC coordination).<sup>151</sup>



In addition to the parent diborene (HB=BH),<sup>143</sup> carbene-stabilized diboron complexes have also been utilized in the synthesis of a variety of diborene derivatives (including the electron-poor cyano-substituted diborene,<sup>152</sup> *trans*-1-(2-propenyloxy)-2-hydrodiborene,<sup>153</sup> and diboryldiborenes<sup>154</sup>). Notably, SNHC<sup>Dipp,Mes</sup>-stabilized diboron complex (**102**) has been shown to undergo stepwise hydroboration reactions with catecholborane to give hydro(boryl)-diborene (**103**) and 2,3-dihydrodiborene (**104**) (Scheme 26).<sup>155</sup> MeOTf-mediated hydride (H<sup>−</sup>) abstraction from **104** affords a unique planar

**Scheme 25.** Activation of Small Molecules by Carbene-Diboron Complexes (L: = NHC<sup>Dipp</sup>; Dipp = 2,6-Diisopropylphenyl, Dep = 2,6-Diethylphenyl)



tetraborane cation (**105**), wherein a hydrogen atom serves as a bridge between the two boron atoms (Scheme 26). Both the solid-state structural data and DFT computations support its “protonated diborene” structural feature.

Braunschweig also utilized diboracumulene (**81**) in the synthesis of boron-containing aromatic heterocycles. While reacting with propyne to produce a CAAC-based 1,3-diborene (**106**), which exists as a triplet biradical with the unpaired electrons residing at the CAAC ligands, **81** may also react with acetylene to give a neutral CAAC-based 1,4-diborabenzene (**107**) via [2 + 2 + 2]-cycloaddition of two acetylene molecules to the B<sub>2</sub> core in **81** (Scheme 27).<sup>156</sup> **107** was further utilized as a 6 $\pi$ -aromatic analogue of benzene to prepare a series of half-sandwich complexes of transition metals (**108–110**) (Scheme 27).<sup>157</sup> Remarkably, unlike that of arene complexes of transition metals, the redox processes of these 1,4-

**Scheme 26. Synthesis of 103–105 (Dipp = 2,6-Diisopropylphenyl, Mes = Mesityl, cat = Catechol)**

**Scheme 27. Synthesis of 106–110 (Dipp = 2,6-Diisopropylphenyl)**



diborabenzene-transition metal complexes are largely ligand-based.

**■ CARBENE-STABILIZED E<sub>N</sub>(0) (N = 3, 4, 8, 12) SPECIES**


Carbene-stabilized triatomic molecules are rare. Grützmacher synthesized NHC-based P<sub>3</sub><sup>+</sup> (111) and P<sub>2</sub>As<sup>+</sup> (112) cations in 2014 by allowing the NHC-stabilized parent phosphinidene (PH) to react with PCl<sub>3</sub> and AsCl<sub>3</sub>, respectively, in the presence of DABCO (1,4-diazabicyclo[2.2.2]octane) (Scheme 28).<sup>158</sup> Subsequent magnesium reduction of 111 and 112 gave NHC-stabilized P<sub>3</sub><sup>•</sup> (113) and P<sub>2</sub>As<sup>•</sup> (114) radicals, respectively (Scheme 28).<sup>158</sup>

The Mulliken spin densities [experimental (theoretical)] are largely located on the central P atom of 113 [64.4% (65.6%)] and the As atom of 114 [80.4% (67.3%)]. The P–C bonds of 113 [1.766(2) Å, av] compare well to that of NHC<sup>Dipp</sup>-stabilized P<sub>2</sub>(0) (59)<sup>113</sup> [1.7504(17) Å]. In addition, the P–P bond distances of 113 [2.144(1) Å, av] are comparable to those of 111 [2.094(1) Å, av] and that of 59 [2.2052(10) Å]. When being combined with the activated alkyne MeOOC≡COOMe, 113 may serve as a “P<sub>3</sub>” transfer agent to give 115 (Scheme 28).<sup>158</sup> This conversion involves cycloaddition between the alkyne unit and the P<sub>3</sub> species, along with subsequent single electron transfer.

After obtaining a series of CAAC-stabilized monatomic<sup>44,61</sup> and diatomic silicon(0) species,<sup>75,101</sup> Roesky synthesized a

**Scheme 28. Synthesis of 111–115**


<sup>Me</sup>CAAC-stabilized triatomic silicon(0) complex (116) in 2016, via potassium graphite reduction of <sup>Me</sup>CAAC:SiCl<sub>4</sub> at extremely low temperature (Scheme 29).<sup>159</sup> It was noted that

**Scheme 29. Synthesis of CAAC-Stabilized Si<sub>3</sub>(0) Ring (116)**


116 is acquired exclusively when the reduction of <sup>Me</sup>CAAC:SiCl<sub>4</sub> proceeds slowly under an argon atmosphere. Otherwise, the reaction would give a mixture of both 116 and <sup>Me</sup>CAAC-stabilized Si<sub>2</sub>(0).<sup>101</sup> Further investigation of the possible conversion between CAAC-stabilized Si<sub>2</sub>(0) and Si<sub>3</sub>(0) species may provide insight for accessing larger silicon clusters.

X-ray structural analysis<sup>159</sup> (Figure 12) shows that 116 contains a three-membered silicon ring, with three silicon–





Figure 12. Molecular structure of 116.

silicon single bonds (2.389 Å, av). Each three-coordinate silicon atom, bearing an electron pair, adopts the trigonal pyramidal geometry. The <sup>29</sup>Si NMR resonance of 116 (+7.20 ppm) is upfield-shifted compared to that of CAAC-stabilized mono silicon(0) atom (14) (+66.71 ppm)<sup>44</sup> and diatomic

silicon (**29**) (+254.60 ppm).<sup>75</sup> The shortened (ca. 0.08 Å) Si–C<sub>CAAC</sub> bonds and elongated (ca. 0.08 Å) N–C<sub>CAAC</sub> bonds of **116** (compared to those of the CAAC:SiCl<sub>4</sub> precursor) reveal significant  $\pi$  back-donation from each silicon(0) atom to the carbene carbon atom.<sup>159</sup> This  $\pi$  back-donation plays a key role in stabilizing the Si<sub>3</sub>(0) core. Notably, the ligand-exchange reaction (CAAC)<sub>3</sub>Si<sub>3</sub> + 3NHC<sup>Ph</sup> → (NHC<sup>Ph</sup>)<sub>3</sub>Si<sub>3</sub> + 3CAAC is endergonic ( $\Delta G = 20.9$  kcal mol<sup>-1</sup>). At present, the NHC-based Si<sub>3</sub>(0) complex remains elusive.

Kinjo recently synthesized a zwitterionic boraalkenyl boronium (**118**) by the reaction of 1,2-azaborole-derived CAAC-complexed B<sub>2</sub>Br<sub>4</sub> (**117**) with 4 equiv of PMe<sub>3</sub> and subsequently with 5 equiv of potassium-graphite (Scheme 30).

### Scheme 30. Synthesis of **118**–**120** (Dipp = 2,6-Diisopropylphenyl)



Further reduction of **118** by potassium-graphite (8 equiv) gave (CAAC and phosphine)-based neutral allenic diborene (**119**) (Scheme 30).<sup>160</sup> Reaction of **119** with **117** (in a 3:1 ratio) yielded a (CAAC and phosphine)-stabilized tetraatomic boron(0) species (**120**) (Scheme 30).<sup>161</sup>

The solid-state structure of compound **120** reveals a distinctive planar four-membered boron ring (Figure 13).<sup>161</sup>





Figure 13. Molecular structure of **120**.

The bridging B1 atom is two-coordinate (bonding only to B2 and B4), whereas the bridging B3 atom is four-coordinate (bonding to B2 and B4, in addition to two PMe<sub>3</sub> ligands). The three-coordinate B2 and B4 atoms bond to the carbene carbon atoms. While the 1.564(3) Å B1–B2 and 1.554(3) Å B1–B4 bonds are double bonds with an identical WBI value of 1.31, the B2–B3 [1.833(3) Å] and B3–B4 [1.849(3) Å] bonds are single bonds with WBI values of 0.69 and 0.68, respectively. The 1.42 WBI values for both B2–C1 [1.456(3) Å] and B4–C2 [1.454(3) Å] bonds reveal the partially multiple bond nature of the boron–carbon bonds in **120**. Computations, coupled with X-ray structural data, indicate that the partial delocalization of electrons in **120** occurs not only in the  $\sigma$ -

framework of the B<sub>4</sub> ring but also in the conjugated  $\pi$ -system over the C1–B2–B1–B4–C2 fragment.

Carbene-mediated activation of white phosphorus (P<sub>4</sub>) has been confirmed to be an effective method to access various carbene-stabilized P<sub>n</sub> clusters.<sup>125,162–165</sup> Bertrand has suggested that the electronic and steric properties of carbenes, as well as the reaction stoichiometry, may impact the formation of carbene-complexed P<sub>n</sub> clusters. While the 4:1 reaction of <sup>Cy</sup>CAAC (**69**) with P<sub>4</sub> in Et<sub>2</sub>O gave both the <sup>Cy</sup>CAAC-stabilized pyramidal P<sub>4</sub> cluster (**70**) and P<sub>2</sub> (**71**) (Scheme 20),<sup>125</sup> the corresponding 1:2 reaction of **69** with P<sub>4</sub> in benzene/toluene mixed solvent afforded the <sup>Cy</sup>CAAC-stabilized P<sub>8</sub> cluster (**122**) (Scheme 31).<sup>163</sup> Interestingly, CAAC-

### Scheme 31. Synthesis of **121**–**123** (R = 2,6-Diisopropylphenyl)



stabilized isomeric P<sub>4</sub> chains [**121a** (*E* isomer, major product) and **121b** (*Z* isomer)] were obtained by employing a more sterically demanding CAAC ligand (Scheme 31).<sup>162</sup> Dimerization of the P<sub>4</sub> core as shown in **121b** via [2 + 2] cycloaddition would give the P<sub>8</sub> cluster as observed in **122**. Notably, carbonyl-modified electrophilic carbenes have also been employed to stabilize the P<sub>8</sub> cluster by the Bertrand and Hudnall laboratories.<sup>163,165</sup> NHCs and CAACs have often demonstrated distinct reactivities. Indeed, the 2:1 reaction of SNHC<sup>Dipp</sup> with P<sub>4</sub> gave a carbene-P<sub>12</sub> complex (**123**), which presently represents the largest elemental cluster stabilized by carbenes (Scheme 31).<sup>164</sup> Both (*E*)-tetraphosphatriene and triphosphirene species have been proposed as remarkable intermediates in the formation of **123**.

X-ray structural analysis (Figure 14) shows that the  $P_8$  core in **122** consists of a central butterfly  $P_4$  ring and four terminal




Figure 14. Molecular structures of **122** and **123**.

P atoms. Each terminal phosphorus atom is bridged between one P atom of the  $P_4$  ring and a  $^{\text{C}}\text{yCAAC}$  ligand.<sup>163</sup> The polycyclic structure of the  $P_{12}$  core in **123** (Figure 14) involves one six-membered ring, three five-membered rings, and two three-membered rings.<sup>164</sup> All of the P–P bonds in **122** (between 2.198 and 2.257 Å) and **123** (between 2.176 and 2.233 Å) are single bonds. The P– $C_{\text{CAAC}}$  bonds in **122** (1.737 Å) are marginally shorter than those for **123** (1.756 and 1.777 Å) due to the stronger electrophilicity of CAACs (than NHCs). Formation of diverse carbene-stabilized  $P_n$  clusters ( $n = 2, 3, 4, 8,$  and  $12$ ) are unusual, which may be due to the unique catenation capability of phosphorus. Notably, carbene stabilization of (large-sized) elemental clusters of p-block metals are challenging, which may be ascribed to the relatively weak  $C_{\text{carbene}}\text{--}E$  bonds ( $E = \text{p-block metals}$ ).

## SUMMARY AND PERSPECTIVE

The chemistry of carbene-stabilized zero-oxidation state main group species is fascinating. Owing to the distinct electronic and steric properties of N-heterocyclic carbenes (NHCs) and cyclic (alkyl)(amino)carbenes (CAACs), the  $E_n(0)$  cores stabilized by these two types of carbene ligands usually exhibit not only different structural and bonding motifs but also contrasting reactivities. The highly electron-rich  $E_n(0)$  cores grant these carbene-stabilized zerovalent main group complexes unusual utilities in coordination chemistry and small molecule activations. For instance, in contrast to the carbene and silylene ligands, which may involve one dative bond with the Lewis acidic species, carbene-complexed  $E(0)$  cores ( $E = \text{C}$  and  $\text{Si}$ ) have exhibited the unique capabilities of binding to two metal centers. The carbene-stabilized  $E_2(0)$  species ( $E = \text{Si}$  and  $\text{Ge}$ ) bearing a  $E\text{=E}$  double bond and two E-based lone pairs have shown different bonding modes toward Lewis acidic species from alkenes. In addition, the carbene-stabilized  $E(0)$  ( $E = \text{Si}$  and  $\text{Ge}$ ) and  $E_2(0)$  ( $E = \text{B}, \text{Si},$  and  $\text{P}$ ) species have

been employed to activate a series of small molecules such as gaseous molecules ( $\text{O}_2, \text{H}_2, \text{CO}, \text{CO}_2, \text{N}_2\text{O}$ ) and elemental chalcogens (sulfur, selenium, and tellurium).

Despite its rapid development, there remain many fascinating challenges in this field. For example, while CDC or silylene-based bimetallics have been obtained, such complexes remain rare. This is largely due to the second latent lone pair of the  $E(0)$  centers in these complexes. The design and synthesis of new  $E(0)$ -containing carbene complexes that possess two potent  $E(0)$ -based donating sites would substantially extend their applications in transition metal coordination chemistry and catalysis. Moreover, stabilization of the highly reactive  $E_2(0)$  cores usually require sterically demanding carbene ligands. However, the steric bulk of the carbene ligands limits the access of many transition metal species. Tuning the steric properties of carbene ligands may enhance the utility of carbene-stabilized  $E_2(0)$  species in transition metal coordination chemistry. Given the seminal discoveries detailed herein, many more wonders in this ascendent field await the imaginative chemist.

## AUTHOR INFORMATION

### Corresponding Author

Gregory H. Robinson – Department of Chemistry, The University of Georgia, Athens, Georgia 30602-2556, United States; [orcid.org/0000-0002-2260-3019](https://orcid.org/0000-0002-2260-3019); Email: [robinson@uga.edu](mailto:robinson@uga.edu)

### Author

Yuzhong Wang – Department of Chemistry, The University of Georgia, Athens, Georgia 30602-2556, United States; [orcid.org/0000-0003-3557-5085](https://orcid.org/0000-0003-3557-5085)

Complete contact information is available at: <https://pubs.acs.org/10.1021/jacs.2c13574>

### Notes

The authors declare no competing financial interest.

## ACKNOWLEDGMENTS

We are grateful to the National Science Foundation for support: CHE-1855641 and CHE-2153978 to G.H.R. and Y.W.

## REFERENCES

- Igau, A.; Grützmacher, H.; Baceiredo, A.; Bertrand, G. Analogous  $\alpha, \alpha'$ -Bis-Carbenoid, Triply Bonded Species: Synthesis of a Stable  $\lambda^3$ -Phosphinocarbene- $\lambda^5$ -Phosphaacetylene. *J. Am. Chem. Soc.* **1988**, *110*, 6463–6466.
- Igau, A.; Baceiredo, A.; Trinquier, G.; Bertrand, G. Bis-(diisopropylamino)phosphinotrimethylsilylcarbene: A Stable Carbene Nucleophile. *Angew. Chem., Int. Ed.* **1989**, *28*, 621–622.
- Nelson, D. J.; Praetorius, J. M.; Crudden, C. M. Synthesis, Activation, and Decomposition of N-Heterocyclic Carbene-Containing Complexes. In *N-Heterocyclic Carbenes: From Laboratory Curiosities to Efficient Synthetic Tools*, 2nd ed.; Díez-González, S., Ed.; The Royal Society of Chemistry: Cambridge, 2017; pp 46–98.
- Arduengo, A. J., III; Harlow, R. L.; Kline, M. A Stable Crystalline Carbene. *J. Am. Chem. Soc.* **1991**, *113*, 361–363.
- Petronilho, A.; Albrecht, M. Non-Classical N-Heterocyclic Carbene Complexes. In *N-Heterocyclic Carbenes: From Laboratory Curiosities to Efficient Synthetic Tools*, 2nd ed.; Díez-González, S., Ed.; The Royal Society of Chemistry: Cambridge, 2017; pp 99–119.



- (6) Soleilhavoup, M.; Bertrand, G. Cyclic (Alkyl)(Amino)Carbenes (CAACs): Stable Carbenes on the Rise. *Acc. Chem. Res.* **2015**, *48*, 256–266.
- (7) Melaimi, M.; Jazzar, R.; Soleilhavoup, M.; Bertrand, G. Cyclic (Alkyl)(amino)carbenes (CAACs): Recent Developments. *Angew. Chem., Int. Ed.* **2017**, *56*, 10046–10068.
- (8) Kushvaha, S. K.; Mishra, A.; Roesky, H. W.; Mondal, K. C. Recent Advances in the Domain of Cyclic (Alkyl)(Amino) Carbenes. *Chem. - Asian J.* **2022**, *17*, e202101301.
- (9) Lavallo, V.; Canac, Y.; Prasang, C.; Donnadiu, B.; Bertrand, G. Stable Cyclic (Alkyl)(Amino)Carbenes as Rigid or Flexible, Bulky, Electron-Rich Ligands for Transition-Metal Catalysts: A Quaternary Carbon Atom Makes the Difference. *Angew. Chem., Int. Ed.* **2005**, *44*, 5705–5709.
- (10) Tukov, A. A.; Normand, A. T.; Nechaev, M. S. N-Heterocyclic Carbenes Bearing Two, One and No Nitrogen Atoms at the Ylidene Carbon: Insight from Theoretical Calculations. *Dalton Trans.* **2009**, 7015–7028.
- (11) Lavallo, V.; Canac, Y.; Donnadiu, B.; Schoeller, W. W.; Bertrand, G. CO Fixation to Stable Acyclic and Cyclic Alkyl Amino Carbenes: Stable Amino Ketenes with a Small HOMO-LUMO Gap. *Angew. Chem., Int. Ed.* **2006**, *45*, 3488–3491.
- (12) Back, O.; Henry-Ellinger, M.; Martin, C. D.; Martin, D.; Bertrand, G. <sup>31</sup>P NMR Chemical Shifts of Carbene-Phosphinidene Adducts as an Indicator of the  $\pi$ -Accepting Properties of Carbenes. *Angew. Chem., Int. Ed.* **2013**, *52*, 2939–2943.
- (13) Frey, G. D.; Lavallo, V.; Donnadiu, B.; Schoeller, W. W.; Bertrand, G. Facile Splitting of Hydrogen and Ammonia by Nucleophilic Activation at a Single Carbon Center. *Science* **2007**, *316*, 439–441.
- (14) *N-Heterocyclic Carbenes in Synthesis*; Nolan, S. P., Ed.; Wiley-VCH: Weinheim, 2006.
- (15) Nesterov, V.; Reiter, D.; Bag, P.; Frisch, P.; Holzner, R.; Porzelt, A.; Inoue, S. NHCs in Main Group Chemistry. *Chem. Rev.* **2018**, *118*, 9678–9842.
- (16) Wang, Y.; Robinson, G. H. Carbene-Stabilized Main Group Diatomic Allotropes. *Dalton Trans.* **2012**, *41*, 337–345.
- (17) Wang, Y.; Robinson, G. H. N-Heterocyclic Carbene—Main-Group Chemistry: A Rapidly Evolving Field. *Inorg. Chem.* **2014**, *53*, 11815–11832.
- (18) Arrowsmith, M.; Braunschweig, H.; Stennett, T. E. Formation and Reactivity of Electron-Precise B–B Single and Multiple Bonds. *Angew. Chem., Int. Ed.* **2017**, *56*, 96–115.
- (19) Martin, D.; Melaimi, M.; Soleilhavoup, M.; Bertrand, G. A Brief Survey of Our Contribution to Stable Carbene Chemistry. *Organometallics* **2011**, *30*, 5304–5313.
- (20) Dyker, C. A.; Bertrand, G. Soluble Allotropes of Main-Group Elements. *Science* **2008**, *321*, 1050–1051.
- (21) Majhi, P. K.; Sasamori, T. Tetrylones: An Intriguing Class of Monoatomic Zero-Valent Group 14 Compounds. *Chem. - Eur. J.* **2018**, *24*, 9441–9455.
- (22) Yao, S.; Xiong, Y.; Driess, M. A New Area in Main-Group Chemistry: Zerovalent Monoatomic Silicon Compounds and Their Analogues. *Acc. Chem. Res.* **2017**, *50*, 2026–2037.
- (23) Yao, S.; Xiong, Y.; Saddington, A.; Driess, M. Entering New Chemical Space with Isolable Complexes of Single, Zero-Valent Silicon and Germanium Atoms. *Chem. Commun.* **2021**, *57*, 10139–10153.
- (24) Ota, K.; Kinjo, R. Zero-Valent Species of Group 13–15 Elements. *Chem.* **2022**, *8*, 340–350.
- (25) Ramirez, F.; Desai, N. B.; Hansen, B.; Mckelvie, N. Hexaphenylcarbodi-phosphorane, (C<sub>6</sub>H<sub>5</sub>)<sub>3</sub>PCP(C<sub>6</sub>H<sub>5</sub>)<sub>3</sub>. *J. Am. Chem. Soc.* **1961**, *83*, 3539–3540.
- (26) Tonner, R.; Öxler, F.; Neumüller, B.; Petz, W.; Frenking, G. Carbodiphosphoranes: The Chemistry of Divalent Carbon(0). *Angew. Chem., Int. Ed.* **2006**, *45*, 8038–8042.
- (27) Tonner, R.; Frenking, G. C(NHC)<sub>2</sub>: Divalent Carbon(0) Compounds with N-Heterocyclic Carbene Ligands-Theoretical Evidence for a Class of Molecules with Promising Chemical Properties. *Angew. Chem., Int. Ed.* **2007**, *46*, 8695–8698.
- (28) Tonner, R.; Frenking, G. Divalent Carbon(0) Chemistry, Part 1: Parent Compounds. *Chem. - Eur. J.* **2008**, *14*, 3260–3272.
- (29) Dyker, C. A.; Lavallo, V.; Donnadiu, B.; Bertrand, G. Synthesis of an Extremely Bent Acyclic Allene (A “Carbodicarbene”): A Strong Donor Ligand. *Angew. Chem., Int. Ed.* **2008**, *47*, 3206–3209.
- (30) Klein, S.; Tonner, R.; Frenking, G. Carbodicarbenes and Related Divalent Carbon(0) Compounds. *Chem. - Eur. J.* **2010**, *16*, 10160–10170.
- (31) Fürstner, A.; Alcarazo, M.; Goddard, R.; Lehmann, C. W. Coordination Chemistry of Ene-1,1-diamines and a Prototype “Carbodicarbene”. *Angew. Chem., Int. Ed.* **2008**, *47*, 3210–3214.
- (32) Bowden, F. L.; Giles, R. The Coordination Chemistry of Allenes. *Coord. Chem. Rev.* **1976**, *20*, 81–106.
- (33) Zhao, L.; Chai, C.; Petz, W.; Frenking, G. Carbones and Carbon Atom as Ligands in Transition Metal Complexes. *Molecules* **2020**, *25*, 4943.
- (34) Au-Yeung, K.-C.; Xiao, D.; Shih, W.-C.; Yang, H.-W.; Wen, Y.-S.; Yap, G. P. A.; Chen, W.-C.; Zhao, L.; Ong, T.-G. Carbodicarbene: geminal-Bimetallic Coordination in Selective Manner. *Chem. - Eur. J.* **2020**, *26*, 17350–17355.
- (35) Chen, W.-C.; Shih, W.-C.; Jurca, T.; Zhao, L.; Andrada, D. M.; Peng, C.-J.; Chang, C.-C.; Liu, S.-K.; Wang, Y.-P.; Wen, Y.-S.; Yap, G. P. A.; Hsu, C.-P.; Frenking, G.; Ong, T.-G. Carbodicarbenes: Unexpected  $\pi$ -Accepting Ability during Reactivity with Small Molecules. *J. Am. Chem. Soc.* **2017**, *139*, 12830–12836.
- (36) Chen, W.-C.; Shen, J.-S.; Jurca, T.; Peng, C.-J.; Lin, Y.-H.; Wang, Y.-P.; Shih, W.-C.; Yap, G. P. A.; Ong, T.-G. Expanding the Ligand Framework Diversity of Carbodicarbenes and Direct Detection of Boron Activation in the Methylation of Amines with CO<sub>2</sub>. *Angew. Chem., Int. Ed.* **2015**, *54*, 15207–15212.
- (37) Liu, S.-K.; Shih, W.-C.; Chen, W.-C.; Ong, T.-G. Carbodicarbenes and their Captodative Behavior in Catalysis. *ChemCatChem* **2018**, *10*, 1483–1498.
- (38) Wang, Y.; Quillian, B.; Wei, P.; Wannere, C. S.; Xie, Y.; King, R. B.; Schaefer, H. F., III; Schleyer, P. v. R.; Robinson, G. H. A Stable Neutral Diborene Containing a B=B Double Bond. *J. Am. Chem. Soc.* **2007**, *129*, 12412–12413.
- (39) Chen, W.-C.; Lee, C.-Y.; Lin, B.-C.; Hsu, Y.-C.; Shen, J.-S.; Hsu, C.-P.; Yap, G. P. A.; Ong, T.-G. The Elusive Three-Coordinate Dicationic Hydrido Boron Complex. *J. Am. Chem. Soc.* **2014**, *136*, 914–917.
- (40) Walley, J. E.; Warring, L. S.; Wang, G.; Dickie, D. A.; Pan, S.; Frenking, G.; Gilliard, R. J., Jr. Carbodicarbene Bismaalkene Cations: Unravelling the Complexities of Carbene versus Carbene in Heavy Pnictogen Chemistry. *Angew. Chem., Int. Ed.* **2021**, *60*, 6682–6690.
- (41) Chan, S.-C.; Ang, Z. Z.; Gupta, P.; Ganguly, R.; Li, Y.; Ye, S.; England, J. Carbodicarbene Ligand Redox Noninnocence in Highly Oxidized Chromium and Cobalt Complexes. *Inorg. Chem.* **2020**, *59*, 4118–4128.
- (42) Takagi, N.; Shimizu, T.; Frenking, G. Divalent Silicon(0) Compounds. *Chem. - Eur. J.* **2009**, *15*, 3448–3456.
- (43) Ishida, S.; Iwamoto, T.; Kabuto, C.; Kira, M. A Stable Silicon-Based Allene Analogue with a Formally sp-Hybridized Silicon Atom. *Nature* **2003**, *421*, 725–727.
- (44) Mondal, K. C.; Roesky, H. W.; Schwarzer, M. C.; Frenking, G.; Niepotter, B.; Wolf, H.; Herbst-Irmer, R.; Stalke, D. A Stable Singlet Biradicaloid Siladicarbene: (L)<sub>2</sub>Si. *Angew. Chem., Int. Ed.* **2013**, *52*, 2963–2967.
- (45) Mondal, K. C.; Roesky, H. W.; Schwarzer, M. C.; Frenking, G.; Tkach, I.; Wolf, H.; Kratzert, D.; Herbst-Irmer, R.; Niepotter, B.; Stalke, D. Conversion of a Singlet Silylene to a Stable Biradical. *Angew. Chem., Int. Ed.* **2013**, *52*, 1801–1805.
- (46) Lee, V. Y.; Sekiguchi, A. *Organometallic Compounds of Low-Coordinate Si, Ge, Sn, and Pb: From Phantom Species to Stable Compounds*; Wiley: Chichester, 2010.
- (47) Li, Y.; Chan, Y.-C.; Li, Y.; Purushothaman, I.; De, S.; Parameswaran, P.; So, C.-W. Synthesis of a Bent 2-Silaallene with a

Perturbed Electronic Structure from a Cyclic Alkyl(amino) Carbene-Diiodosilylene. *Inorg. Chem.* **2016**, *55*, 9091–9098.

(48) Xiong, Y.; Yao, S.; Tan, G.; Inoue, S.; Driess, M. A Cyclic Germadecarbene ("Germylone") from Germyliumylidene. *J. Am. Chem. Soc.* **2013**, *135*, 5004–5007.

(49) Sidiropoulos, A.; Jones, C.; Stasch, A.; Klein, S.; Frenking, G. N-Heterocyclic Carbene Stabilized Digermanium(0). *Angew. Chem., Int. Ed.* **2009**, *48*, 9701–9704.

(50) Li, Y.; Mondal, K. C.; Roesky, H. W.; Zhu, H.; Stollberg, P.; Herbst-Irmer, R.; Stalke, D.; Andradá, D. M. Acyclic Germylones: Congeners of Allenes with a Central Germanium Atom. *J. Am. Chem. Soc.* **2013**, *135*, 12422–12428.

(51) Xiong, Y.; Yao, S.; Inoue, S.; Epping, J. D.; Driess, M. A Cyclic Silylone ("Siladecarbene") with an Electron-Rich Silicon(0). *Atom Angew. Chem. Int. Ed.* **2013**, *52*, 7147–7150.

(52) Wu, M.; He, Y.; Zhang, L.; Wei, R.; Wang, D.; Liu, J.; Liu, L. L.; Tan, G. An Acyclic Silylone Stabilized by Mesoionic Carbene. *Eur. J. Inorg. Chem.* **2022**, *2022*, e202200413.

(53) Barnett, C.; Cole, M. L.; Harper, J. B. The Core Difference Between a Mesoionic and a Normal N-Heterocyclic Carbene. *ACS Omega* **2022**, *7*, 34657–34664.

(54) Su, B.; Ganguly, R.; Li, Y.; Kinjo, R. Isolation of an Imino-N-Heterocyclic Carbene/Germanium(0) Adduct: A Mesoionic Germylene Equivalent. *Angew. Chem., Int. Ed.* **2014**, *53*, 13106–13109.

(55) Nguyen, M. T.; Gusev, D.; Dmitrienko, A.; Gabidullin, B. M.; Spasyuk, D.; Pilkington, M.; Nikonov, G. I. Ge(0) Compound Stabilized by a Diimino-Carbene Ligand: Synthesis and Ambiphilic Reactivity. *J. Am. Chem. Soc.* **2020**, *142*, 5852–5861.

(56) Mandal, D.; Dhara, D.; Maiti, A.; Klemmer, L.; Huch, V.; Zimmer, M.; Rzepa, H. S.; Scheschke, D.; Jana, A. Mono- and Dicoordinate Germanium(0) as a Four-Electron Donor. *Chem. - Eur. J.* **2018**, *24*, 2873–2878.

(57) Burchert, A.; Yao, S.; Müller, R.; Schattenberg, C.; Xiong, Y.; Kaupp, M.; Driess, M. An Isolable Silicon Dicarboxylate Complex from Carbon Dioxide Activation with a Silylone. *Angew. Chem., Int. Ed.* **2017**, *56*, 1894–1897.

(58) Xiong, Y.; Yao, S. L.; Müller, R.; Kaupp, M.; Driess, M. From Silylone to an Isolable Monomeric Silicon Disulfide Complex. *Angew. Chem., Int. Ed.* **2015**, *54*, 10254–10257.

(59) Burchert, A.; Müller, R.; Yao, S. L.; Schattenberg, C.; Xiong, Y.; Kaupp, M.; Driess, M. Taming Silicon Congeners of CO and CO<sub>2</sub>: Synthesis of Monomeric Si<sup>II</sup> and Si<sup>IV</sup> Chalcogenide Complexes. *Angew. Chem., Int. Ed.* **2017**, *56*, 6298–6301.

(60) Xiong, Y.; Yao, S. L.; Karni, M.; Kostenko, A.; Burchert, A.; Apeloig, Y.; Driess, M. Heavier Congeners of CO and CO<sub>2</sub> as Ligands: from Zero-Valent Germanium ('germylone') to Isolable Monomeric GeX and GeX<sub>2</sub> Complexes (X = S, Se, Te). *Chem. Sci.* **2016**, *7*, 5462–5469.

(61) Mondal, K. C.; Samuel, P. P.; Tretiakov, M.; Singh, A. P.; Roesky, H. W.; Stueckl, A. C.; Niepoetter, B.; Carl, E.; Wolf, H.; Herbst-Irmer, R.; Stalke, D. Easy Access to Silicon(0) and Silicon(II) Compounds. *Inorg. Chem.* **2013**, *52*, 4736–4743.

(62) Takagi, N.; Shimizu, T.; Frenking, G. Divalent E(0) Compounds (E = Si–Sn). *Chem. - Eur. J.* **2009**, *15*, 8593–8604.

(63) Takagi, N.; Frenking, G. Divalent Pb(0) Compounds. *Theor. Chem. Acc.* **2011**, *129*, 615–623.

(64) Xu, J.; Pan, S.; Yao, S.; Frenking, G.; Driess, M. The Heaviest Bottleable Metallone: Synthesis of a Monatomic, Zero-Valent Lead Complex ("Plumbylone"). *Angew. Chem., Int. Ed.* **2022**, *61*, e202209442.

(65) Chen, M.; Zhang, Z.; Qiao, Z.; Zhao, L.; Mo, Z. An Isolable Bis(Germylene)-Stabilized Plumbylone. *Angew. Chem., Int. Ed.* **2023**, *62*, e202215146.

(66) Couchman, S. A.; Holzmann, N.; Frenking, G.; Wilson, D. J. D.; Dutton, J. L. Beryllium Chemistry the Safe Way: A Theoretical Evaluation of Low Oxidation State Beryllium Compounds. *Dalton Trans.* **2013**, *42*, 11375–11384.

(67) Arrowsmith, M.; Braunschweig, H.; Celik, M. A.; Dellermann, T.; Dewhurst, R. D.; Ewing, W. C.; Hammond, K.; Kramer, T.;

Krummenacher, I.; Mies, J.; Radacki, K.; Schuster, J. K. Neutral Zero-Valent s-Block Complexes with Strong Multiple Bonding. *Nat. Chem.* **2016**, *8*, 890–894.

(68) West, R.; Fink, M. J.; Michl, J. Tetramesityldisilene, A Stable Compound Containing a Silicon–Silicon Double Bond. *Science* **1981**, *214*, 1343–1344.

(69) Sekiguchi, A.; Kinjo, R.; Ichinohe, M. A Stable Compound Containing a Silicon–Silicon Triple Bond. *Science* **2004**, *305*, 1755–1757.

(70) Van Zee, R. J.; Ferrante, R. F.; Weltner, W., Jr. Diatomic Silicon, Silyl, and Oxosilyl (HSiO) Molecules: ESR at 4 K. *J. Chem. Phys.* **1985**, *83*, 6181–6187.

(71) Wang, Y.; Xie, Y.; Wei, P.; King, R. B.; Schaefer, H. F., III; Schleyer, P. v. R.; Robinson, G. H. A Stable Silicon(0) Compound with a Si=Si Double Bond. *Science* **2008**, *321*, 1069–1071.

(72) Wagner, J. P.; Schreiner, P. R. London Dispersion Decisively Contributes to the Thermodynamic Stability of Bulky NHC-Coordinated Main Group Compounds. *J. Chem. Theory Comput.* **2016**, *12*, 231–237.

(73) Arz, M. I.; Straßmann, M.; Meyer, A.; Schnakenburg, G.; Schiemann, O.; Filippou, A. C. One-Electron Oxidation of a Disilicon(0) Compound: An Experimental and Theoretical Study of [Si<sub>2</sub>]<sup>+</sup> Trapped by N-Heterocyclic Carbenes. *Chem. - Eur. J.* **2015**, *21*, 12509–12516.

(74) Ghadwal, R. S.; Roesky, H. W.; Merkel, S.; Henn, J.; Stalke, D. Lewis Base Stabilized Dichlorosilylene. *Angew. Chem., Int. Ed.* **2009**, *48*, 5683–5686.

(75) Mondal, K. C.; Samuel, P. P.; Roesky, H. W.; Aysin, R. R.; Leites, L. A.; Neudeck, S.; Luebben, J.; Dittrich, B.; Holzmann, N.; Hermann, M.; Frenking, G. One-Electron-Mediated Rearrangements of 2,3-Disiladecarbene. *J. Am. Chem. Soc.* **2014**, *136*, 8919–8922.

(76) Nimlos, M. R.; Harding, L. B.; Ellison, G. B. The Electronic States of Si<sub>2</sub> and Si<sub>2</sub><sup>−</sup> as Revealed by Photoelectron Spectroscopy. *J. Chem. Phys.* **1987**, *87*, 5116–5124.

(77) Wang, Y.; Hickox, H. P.; Xie, Y.; Wei, P.; Blair, S. A.; Johnson, M. K.; Schaefer, H. F., III; Robinson, G. H. A Stable Anionic Dithiolene Radical. *J. Am. Chem. Soc.* **2017**, *139*, 6859–6862.

(78) Wang, Y.; Tope, C. A.; Xie, Y.; Wei, P.; Urbauer, J. L.; Schaefer, H. F., III; Robinson, G. H. Carbene-Stabilized Disilicon as a Silicon-Transfer Agent: Synthesis of a Dianionic Silicon Tris(dithiolene) Complex. *Angew. Chem., Int. Ed.* **2020**, *59*, 8864–8867.

(79) Dielmann, F.; Back, O.; Henry-Ellinger, M.; Jerabek, P.; Frenking, G.; Bertrand, G. A Crystalline Singlet Phosphonitrene: A Nitrogen Atom-Transfer Agent. *Science* **2012**, *337*, 1526–1528.

(80) Wang, Y.; Szilvási, T.; Yao, S.; Driess, M. A Bis(silylene)-Stabilized Diphosphorus Compound and its Reactivity as a Monophosphorus Anion Transfer Reagent. *Nat. Chem.* **2020**, *12*, 801–807.

(81) Robinson, T. P.; Cowley, M. J.; Scheschke, D.; Goicoechea, J. M. Phosphide Delivery to a Cyclotrisilene. *Angew. Chem., Int. Ed.* **2015**, *54*, 683–686.

(82) Koike, T.; Nukazawa, T.; Iwamoto, T. Conformationally Switchable Silylone: Electron Redistribution Accompanied by Ligand Reorientation around a Monatomic Silicon. *J. Am. Chem. Soc.* **2021**, *143*, 14332–14341.

(83) Zhang, X.; Wang, H.; Lan, X.; Mei, Y.; Ruiz, D. A.; Liu, L. L. Transfer of an Aluminium Atom: An Avenue to Aluminium Heterocycles. *CCS Chem.* **2022**, *1*. Just Published.

(84) Chen, M.; Wang, Y.; Xie, Y.; Wei, P.; Gilliard, R. J., Jr.; Schwartz, N. A.; Schaefer, H. F., III; Schleyer, P. v. R.; Robinson, G. H. Dynamic Complexation of Copper(I) Chloride by Carbene-Stabilized Disilicon. *Chem. - Eur. J.* **2014**, *20* (20), 9208–9211.

(85) Cowley, M. J.; Abersfelder, K.; White, A. J. P.; Majumdar, M.; Scheschke, D. Transmetalation Reactions of a Lithium Disilene. *Chem. Commun.* **2012**, *48*, 6595–6597.

(86) Kira, M. Bonding and Structure of Disilenes and Related Unsaturated Group-14 Element Compounds. *Proc. Jpn. Acad., Ser. B* **2012**, *88*, 167–191.

(87) Tsutsui, M.; Courtney, A.  $\sigma$ - $\pi$  Rearrangements of Organotransition Metal Compounds. In *Advances in Organometallic*

Chemistry; Stone, F. G. A., West, R., Eds.; Elsevier: New York, 1977; Vol. 16, pp 241–282.

(88) Arz, M. I.; Straßmann, M.; Geiß, D.; Schnakenburg, G.; Filippou, A. C. Addition of Small Electrophiles to N-Heterocyclic-Carbene-Stabilized Disilicon(0): A Revisit of the Isolobal Concept in Low-Valent Silicon Chemistry. *J. Am. Chem. Soc.* **2016**, *138*, 4589–4600.

(89) Arz, M. I.; Geiß, D.; Straßmann, M.; Schnakenburg, G.; Filippou, A. C. Silicon(I) Chemistry: the NHC-Stabilised Silicon(I) Halides  $\text{Si}_2\text{X}_2(\text{Idipp})_2$  (X = Br, I) and the Disilicon(I)-Iodido Cation  $[\text{Si}_2(\text{I})(\text{Idipp})_2]^+$ . *Chem. Sci.* **2015**, *6*, 6515–6524.

(90) Abraham, M. Y.; Wang, Y.; Xie, Y.; Wei, P.; Schaefer, H. F., III; Schleyer, P. v. R.; Robinson, G. H. Cleavage of Carbene-Stabilized Disilicon. *J. Am. Chem. Soc.* **2011**, *133*, 8874–8876.

(91) Köcher, T.; Kerst, C.; Friedrichs, G.; Temps, F. The Gas-Phase Oxidation of Silyl Radicals by Molecular Oxygen: Kinetics and Mechanisms. In *Silicon Chemistry*; Jutzi, P., Schubert, U., Eds.; Wiley-VCH: Weinheim, 2003; pp 44–57.

(92) Hickox, H. P.; Wang, Y.; Xie, Y.; Chen, M.; Wei, P.; Schaefer, H. F., III; Robinson, G. H. Transition Metal-Mediated Cleavage of a Si=Si Double Bond. *Angew. Chem., Int. Ed.* **2015**, *54*, 10267–10270.

(93) Hickox, H. P.; Wang, Y. Z.; Xie, Y. M.; Wei, P. R.; Schaefer, H. F., III; Robinson, G. H. Push-Pull Stabilization of Parent Monochlorosilylenes. *J. Am. Chem. Soc.* **2016**, *138*, 9799–9802.

(94) *Silicon Chemistry: From the Atom to Extended Systems*; Jutzi, P.; Schubert, U., Eds.; Wiley-VCH: Weinheim, 2003.

(95) Schnöckel, H. IR Spectroscopic Detection of Molecular  $\text{SiO}_2$ . *Angew. Chem., Int. Ed.* **1978**, *17*, 616–617.

(96) Chakraborty, S.; Yanchulova, P.; Thiemens, M. H. Mass-Independent Oxygen Isotopic Partitioning during Gas-Phase  $\text{SiO}_2$  Formation. *Science* **2013**, *342*, 463–466.

(97) Wang, L.-S.; Wu, H.; Desai, S. R.; Fan, J.; Colson, S. D. A Photoelectron Spectroscopic Study of Small Silicon Oxide Clusters:  $\text{SiO}_2$ ,  $\text{Si}_2\text{O}_3$ , and  $\text{Si}_2\text{O}_4$ . *J. Phys. Chem.* **1996**, *100*, 8697–8700.

(98) Wang, Y.; Chen, M.; Xie, Y.; Wei, P.; Schaefer, H. F., III; Schleyer, P. v. R.; Robinson, G. H. Stabilization of Elusive Silicon Oxides. *Nat. Chem.* **2015**, *7*, 509–513.

(99) Wang, Y.; Chen, M.; Xie, Y.; Wei, P.; Schaefer, H. F., III; Robinson, G. H. Stabilization of Silicon-Carbon Mixed Oxides. *J. Am. Chem. Soc.* **2015**, *137*, 8396–8399.

(100) Chan, Y.-C.; Leong, B.-X.; Li, Y.; Yang, M.-C.; Li, Y.; Su, M.-D.; So, C.-W. A Dimeric NHC-Silicon Monotelluride: Synthesis, Isomerization, and Reactivity. *Angew. Chem., Int. Ed.* **2017**, *56*, 11565–11569.

(101) Chandra Mondal, K.; Roy, S.; Dittrich, B.; Maity, B.; Dutta, S.; Koley, D.; Vasa, S. K.; Linser, R.; Dechert, S.; Roesky, H. W. A Soluble Molecular Variant of the Semiconducting Silicondiselenide. *Chem. Sci.* **2015**, *6*, 5230–5234.

(102) Mohapatra, C.; Mondal, K. C.; Samuel, P. P.; Keil, H.; Niepoetter, B.; Herbst-Irmer, R.; Stalke, D.; Dutta, S.; Koley, D.; Roesky, H. W. A Stable Dimer of  $\text{SiS}_2$  Arranged between Two Carbene Molecules. *Chem. - Eur. J.* **2015**, *21*, 12572–12576.

(103) Jones, C.; Sidiropoulos, A.; Holzmann, N.; Frenking, G.; Stasch, A. An N-Heterocyclic Carbene Adduct of Diatomic Tin;  $\text{Sn}=\text{Sn}$ : *Chem. Commun.* **2012**, *48*, 9855–9857.

(104) Jana, A.; Huch, V.; Rzepa, H. S.; Scheschke, D. A Molecular Complex with a Formally Neutral Iron Germanide Motif ( $\text{Fe}_2\text{Ge}_2$ ). *Organometallics* **2015**, *34*, 2130–2133.

(105) Dutton, J. L.; Wilson, D. J. D. Lewis Base Stabilized Dicarbons: Predictions from Theory. *Angew. Chem., Int. Ed.* **2012**, *51*, 1477–1480.

(106) Georgiou, D. C.; Mahmood, I.; Haghghatbin, M. A.; Hogan, C. F.; Dutton, J. L. The Final Fate of NHC Stabilized Dicarbons. *Pure Appl. Chem.* **2017**, *89*, 791–800.

(107) Li, Y.; Mondal, K. C.; Samuel, P. P.; Zhu, H.; Orben, C. M.; Panneerselvam, S.; Dittrich, B.; Schwederski, B.; Kaim, W.; Mondal, T.; Koley, D.; Roesky, H. W.  $\text{C}_4$  Cumulene and the Corresponding Air-Stable Radical Cation and Dication. *Angew. Chem., Int. Ed.* **2014**, *53*, 4168–4172.

(108) Jin, L.; Melaimi, M.; Liu, L.; Bertrand, G. Singlet Carbenes as Mimics for Transition Metals: Synthesis of an Air Stable Organic Mixed Valence Compound  $[\text{M}_2(\text{C}_2)^{+}]$ ; M = Cyclic(Alkyl)(Amino)-Carbene]. *Org. Chem. Front.* **2014**, *1*, 351–354.

(109) Wu, D.; Li, Y.; Ganguly, R.; Kinjo, R. Synthesis and Structural Characterization of a  $\text{C}_4$  Cumulene Including 4-Pyridylidene Units, and its Reactivity Towards Ammonia-Borane. *Chem. Commun.* **2014**, *50*, 12378–12381.

(110) Gorantla, S. M. N. V. T.; Pan, S.; Mondal, K. C.; Frenking, G. Revisiting the Bonding Scenario of Two Donor Ligand Stabilized  $\text{C}_2$  Species. *J. Phys. Chem. A* **2021**, *125*, 291–301.

(111) Leung, T.-F.; Jiang, D.; Wu, M.-C.; Xiao, D.; Ching, W.-M.; Yap, G. P. A.; Yang, T.; Zhao, L.; Ong, T.-G.; Frenking, G. Isolable Dicarbons Stabilized by a Single Phosphine Ligand. *Nat. Chem.* **2021**, *13*, 89–94.

(112) Wu, M.-C.; Liang, Y.-F.; Jurca, T.; Yap, G. P. A.; Leung, T.-F.; Ong, T.-G. Reactive Dicarbons as a Flexible Ligand for Transition-Metal Coordination and Catalysis. *J. Am. Chem. Soc.* **2022**, *144*, 12996–13005.

(113) Wang, Y.; Xie, Y.; Wei, P.; King, R. B.; Schaefer, H. F., III; Schleyer, P. v. R.; Robinson, G. H. Carbene-Stabilized Diphosphorus. *J. Am. Chem. Soc.* **2008**, *130*, 14970–14971.

(114) Abraham, M. Y.; Wang, Y.; Xie, Y.; Wei, P.; Schaefer, H. F., III; Schleyer, P. v. R.; Robinson, G. H. Carbene Stabilization of Diarsenic: From Hypervalency to Allotropy. *Chem.—Eur. J.* **2010**, *16*, 432–435.

(115) Gilliard, R. J., Jr.; Suter, R.; Schrader, E.; Benkő, Z.; Rheingold, A. L.; Grützmacher, H.; Protasiewicz, J. D. Synthesis of  $\text{P}_2\text{C}_2\text{O}_2$  and  $\text{P}_2\text{CO}$  via NHC-Mediated Coupling of the Phosphaethynolate Anion. *Chem. Commun.* **2017**, *53*, 12325–12328.

(116) Weber, L. Phosphaalkenes with Inverse Electron Density. *Eur. J. Inorg. Chem.* **2000**, *2000*, 2425–2441.

(117) Wang, Y.; Xie, Y.; Abraham, M. Y.; Gilliard, R. J., Jr.; Wei, P.; Schaefer, H. F., III; Schleyer, P. v. R.; Robinson, G. H. Carbene-Stabilized Parent Phosphinidene. *Organometallics* **2010**, *29*, 4778–4780.

(118) Wang, Y.; Xie, Y.; Abraham, M. Y.; Wei, P.; Schaefer, H. F., III; Schleyer, P. v. R.; Robinson, G. H. Carbene-Stabilized Diphosphorus: Bidentate Complexation of  $\text{BH}_2^+$ . *Chem. Commun.* **2011**, *47*, 9224–9226.

(119) Wang, Y.; Xie, Y.; Wei, P.; Schaefer, H. F., III; Schleyer, P. v. R.; Robinson, G. H. Splitting Molecular Oxygen en Route to a Stable Molecule Containing Diphosphorus Tetroxide. *J. Am. Chem. Soc.* **2013**, *135* (135), 19139–19142.

(120) Borovik, A. S.; Zinn, P. J.; Zart, M. K. Dioxygen Binding and Activation: Reactive Intermediates. In *Activation of Small Molecules: Organometallic and Bioinorganic Perspectives*; Tolman, W. B., Ed.; Wiley-VCH: Weinheim, 2006; pp 187.

(121) Lohr, L. L., Jr. An ab Initio Characterization of the Gaseous Diphosphorus Oxides  $\text{P}_2\text{O}_x$  (x = 1–5). *J. Phys. Chem.* **1990**, *94*, 1807–1811.

(122) Bauschlicher, C. W., Jr.; Zhou, M.; Andrews, L. A Study of the Products of the Reaction of Phosphorus and Dioxygen. *J. Phys. Chem. A* **2000**, *104*, 3566–3571.

(123) Back, O.; Donnadiou, B.; Parameswaran, P.; Frenking, G.; Bertrand, G. Isolation of Crystalline Carbene-Stabilized  $\text{P}_2$ -Radical Cations and  $\text{P}_2$ -Dications. *Nat. Chem.* **2010**, *2*, 369–373.

(124) Abraham, M. Y.; Wang, Y.; Xie, Y.; Gilliard, R. J., Jr.; Wei, P.; Vaccaro, B. J.; Johnson, M. K.; Schaefer, H. F., III; Schleyer, P. v. R.; Robinson, G. H. Oxidation of Carbene-Stabilized Diarsenic: Diarsene Dications and Diarsenic Radical Cations. *J. Am. Chem. Soc.* **2013**, *135*, 2486–2488.

(125) Back, O.; Kuchenbeiser, G.; Donnadiou, B.; Bertrand, G. Nonmetal-Mediated Fragmentation of  $\text{P}_4$ : Isolation of  $\text{P}_1$  and  $\text{P}_2$  Bis(carbene) Adducts. *Angew. Chem., Int. Ed.* **2009**, *48*, 5530–5533.

(126) Melancon, K. M.; Gildner, M. B.; Hudnall, T. W. Synthesis, Spectroscopic Characterization, and Redox Reactivity of a Cyclic (Alkyl) Amino Carbene-Derived Arsamethine Cyanine Dye. *Chem.—Eur. J.* **2018**, *24*, 9264–9268.



- (127) Kretschmer, R.; Ruiz, D. A.; Moore, C. E.; Rheingold, A. L.; Bertrand, G. One-, Two-, and Three-Electron Reduction of a Cyclic Alkyl(amino)carbene-SbCl<sub>3</sub> Adduct. *Angew. Chem., Int. Ed.* **2014**, *53*, 8176–8179.
- (128) Dobbs, K. D.; Boggs, J. E.; Cowley, A. H. Trends in Structure and Reactivity of Group 15 Alkyls, Alkylidenes, and Alkylidyne. *Chem. Phys. Lett.* **1987**, *141*, 372–375.
- (129) Robinson, G. H. Gallanes, Gallenes, Cyclogallenes, and Gallynes: Organometallic Chemistry About the Gallium–Gallium Bond. *Acc. Chem. Res.* **1999**, *32*, 773–782.
- (130) Wang, Y.; Robinson, G. H. Organometallics of the Group 13 M–M Bond (M = Al, Ga, In) and the Concept of Metalloaromaticity. *Organometallics* **2007**, *26*, 2–11.
- (131) Moezzi, A.; Olmstead, M. M.; Power, P. P. Boron–Boron Double Bonding in the Species [B<sub>2</sub>R<sub>4</sub>]<sup>2-</sup>: Synthesis and Structure of [(Et<sub>2</sub>O)Li]<sub>2</sub>{Mes<sub>2</sub>BB(Mes)Ph}, a Diborane(4) Dianion Analogue of a Substituted Ethylene. *J. Am. Chem. Soc.* **1992**, *114*, 2715–2717.
- (132) Moezzi, A.; Bartlett, R. A.; Power, P. P. Reduction of a Boron–Nitrogen Analogue of 1,3-Butadiene: Proof of a Strong Boron–Boron  $\pi$ -Bond. *Angew. Chem., Int. Ed.* **1992**, *31*, 1082–1083.
- (133) Cheng, T. C.; Headley, L.; Halasa, A. F. Reaction of Sodium Naphthalene with Halobenzenes. *J. Am. Chem. Soc.* **1971**, *93*, 1502–1503.
- (134) Wang, Y.; Quillian, B.; Wei, P.; Xie, Y.; Wannere, C. S.; King, R. B.; Schaefer, H. F., III; Schleyer, P. v. R.; Robinson, G. H. Planar, Twisted, and Trans-Bent: Conformational Flexibility of Neutral Diborenes. *J. Am. Chem. Soc.* **2008**, *130*, 3298–3299.
- (135) Braunschweig, H.; Dewhurst, R. D.; Hammond, K.; Mies, J.; Radacki, K.; Vargas, A. Ambient-Temperature Isolation of a Compound with a Boron–Boron Triple Bond. *Science* **2012**, *336*, 1420–1422.
- (136) Zhou, M.; Tsumori, N.; Li, Z.; Fan, K.; Andrews, L.; Xu, Q. OCBBCO: A Neutral Molecule with Some Boron–Boron Triple Bond Character. *J. Am. Chem. Soc.* **2002**, *124*, 12936–12937.
- (137) Böhnke, J.; Braunschweig, H.; Constantinidis, P.; Dellermann, T.; Ewing, W. C.; Fischer, I.; Hammond, K.; Hupp, F.; Mies, J.; Schmitt, H. C.; Vargas, A. Experimental Assessment of the Strengths of B–B Triple Bonds. *J. Am. Chem. Soc.* **2015**, *137*, 1766–1769.
- (138) Perras, F. A.; Ewing, W. C.; Dellermann, T.; Böhnke, J.; Ullrich, S.; Schäfer, T.; Braunschweig, H.; Bryce, D. L. Spying on the Boron–Boron Triple Bond Using Spin–Spin Coupling Measured from <sup>11</sup>B Solid-State NMR Spectroscopy. *Chem. Sci.* **2015**, *6*, 3378–3382.
- (139) Böhnke, J.; Braunschweig, H.; Ewing, W. C.; Horl, C.; Kramer, T.; Krummenacher, I.; Mies, J.; Vargas, A. Diborabutatriene: An Electron-Deficient Cumulene. *Angew. Chem., Int. Ed.* **2014**, *53*, 9082–9085.
- (140) Power, P. P.  $\pi$ -Bonding and the Lone Pair Effect in Multiple Bonds between Heavier Main Group Elements. *Chem. Rev.* **1999**, *99*, 3463–3503.
- (141) Fischer, R. C.; Power, P. P.  $\pi$ -Bonding and the Lone Pair Effect in Multiple Bonds Involving Heavier Main Group Elements: Developments in the New Millennium. *Chem. Rev.* **2010**, *110*, 3877–3923.
- (142) Böhnke, J.; Braunschweig, H.; Dellermann, T.; Ewing, W. C.; Hammond, K.; Jiménez-Halla, J. O. C.; Kramer, T.; Mies, J. The Synthesis of B<sub>2</sub>(SIDip)<sub>2</sub> and its Reactivity Between the Diboracumulenic and Diborynic Extremes. *Angew. Chem., Int. Ed.* **2015**, *54*, 13801–13805.
- (143) Arrowsmith, M.; Böhnke, J.; Braunschweig, H.; Celik, M. A.; Dellermann, T.; Hammond, K. Uncatalyzed Hydrogenation of First-Row Main Group Multiple Bonds. *Chem. - Eur. J.* **2016**, *22*, 17169–17172.
- (144) Braunschweig, H.; Dellermann, T.; Dewhurst, R. D.; Hupp, B.; Kramer, T.; Mattock, J. D.; Mies, J.; Phukan, A. K.; Steffen, A.; Vargas, A. Strongly Phosphorescent Transition Metal  $\pi$ -Complexes of Boron–Boron Triple Bonds. *J. Am. Chem. Soc.* **2017**, *139*, 4887–4893.
- (145) Braunschweig, H.; Constantinidis, P.; Dellermann, T.; Ewing, W. C.; Fischer, I.; Hess, M.; Knight, F. R.; Rempel, A.; Schneider, C.; Ullrich, S.; Vargas, A.; Woollins, J. D. Highly Strained Heterocycles Constructed from Boron–Boron Multiple Bonds and Heavy Chalcogens. *Angew. Chem., Int. Ed.* **2016**, *55*, 5606–5609.
- (146) Ewing, W. C.; Dellermann, T.; Wong, Y. T. A.; Mattock, J. D.; Vargas, A.; Bryce, D. L.; Dewhurst, R. D.; Braunschweig, H.  $\pi$ -Complexes of Diborynes with Main Group Atoms. *Chem. - Asian J.* **2020**, *15*, 1553–1557.
- (147) Bertermann, R.; Braunschweig, H.; Constantinidis, P.; Dellermann, T.; Dewhurst, R. D.; Ewing, W. C.; Fischer, I.; Kramer, T.; Mies, J.; Phukan, A. K.; Vargas, A. Exclusive  $\pi$  Encapsulation of Light Alkali Metal Cations by a Neutral Molecule. *Angew. Chem., Int. Ed.* **2015**, *54*, 13090–13094.
- (148) Braunschweig, H.; Dellermann, T.; Ewing, W. C.; Kramer, T.; Schneider, C.; Ullrich, S. Reductive Insertion of Elemental Chalcogens into Boron–Boron Multiple Bonds. *Angew. Chem., Int. Ed.* **2015**, *54*, 10271–10275.
- (149) Stoy, A.; Böhnke, J.; Jiménez-Halla, J. O. C.; Dewhurst, R. D.; Thiess, T.; Braunschweig, H. CO<sub>2</sub> Binding and Splitting by Boron–Boron Multiple Bonds. *Angew. Chem., Int. Ed.* **2018**, *57*, 5947–5951.
- (150) Braunschweig, H.; Dellermann, T.; Dewhurst, R. D.; Ewing, W. C.; Hammond, K.; Jiménez-Halla, J. O. C.; Kramer, T.; Krummenacher, I.; Mies, J.; Phukan, A. K.; Vargas, A. Metal-Free Binding and Coupling of Carbon Monoxide at a Boron–Boron Triple Bond. *Nat. Chem.* **2013**, *5*, 1025–1028.
- (151) Böhnke, J.; Arrowsmith, M.; Braunschweig, H. Reactivity Enhancement of a Zerovalent Diboron Compound by Desymmetrization. *J. Am. Chem. Soc.* **2018**, *140*, 10368–10373.
- (152) Böhnke, J.; Braunschweig, H.; Dellermann, T.; Ewing, W. C.; Kramer, T.; Krummenacher, I.; Vargas, A. From an Electron-Rich Bis(boraketimine) to an Electron-Poor Diborene. *Angew. Chem., Int. Ed.* **2015**, *54*, 4469–4473.
- (153) Böhnke, J.; Brückner, T.; Hermann, A.; González-Belman, O. F.; Arrowsmith, M.; Jiménez-Halla, J. O. C.; Braunschweig, H. Single and Double Activation of Acetone by Isolobal B $\equiv$ N and B $\equiv$ B Triple Bonds. *Chem. Sci.* **2018**, *9*, 5354–5359.
- (154) Brückner, T.; Dewhurst, R. D.; Dellermann, T.; Müller, M.; Braunschweig, H. Mild Synthesis of Diboryldiborenes by Diboration of B–B Triple Bonds. *Chem. Sci.* **2019**, *10*, 7375–7378.
- (155) Brückner, T.; Stennett, T. E.; Hess, M.; Braunschweig, H. Single and Double Hydroboration of B–B Triple Bonds and Convergent Routes to a Cationic Tetraborane. *J. Am. Chem. Soc.* **2019**, *141*, 14898–14903.
- (156) Arrowsmith, M.; Böhnke, J.; Braunschweig, H.; Celik, M. A.; Claes, C.; Ewing, W. C.; Krummenacher, I.; Lubitz, K.; Schneider, C. Neutral Diboron Analogues of Archetypal Aromatic Species by Spontaneous Cycloaddition. *Angew. Chem., Int. Ed.* **2016**, *55*, 11271–11275.
- (157) Böhnke, J.; Braunschweig, H.; Jiménez-Halla, J. O. C.; Krummenacher, I.; Stennett, T. E. Half-Sandwich Complexes of an Extremely Electron-Donating, Redox-Active  $\eta^6$ -Diborabenzene Ligand. *J. Am. Chem. Soc.* **2018**, *140*, 848–853.
- (158) Tondreau, A. M.; Benkó, Z.; Harmer, J. R.; Grützmacher, H. Sodium Phosphaethynolate, Na(OCP), as a “P” Transfer Reagent for the Synthesis of N-Heterocyclic Carbene Supported P<sub>3</sub> and PAsP Radicals. *Chem. Sci.* **2014**, *5*, 1545–1554.
- (159) Mondal, K. C.; Roy, S.; Dittrich, B.; Andrada, D. M.; Frenking, G.; Roesky, H. W. A Triatomic Silicon(0) Cluster Stabilized by a Cyclic Alkyl(amino) Carbene. *Angew. Chem., Int. Ed.* **2016**, *55*, 3158–3161.
- (160) Lu, W.; Li, Y.; Ganguly, R.; Kinjo, R. Crystalline Neutral Allenic Diborene. *Angew. Chem., Int. Ed.* **2017**, *56*, 9829–9832.
- (161) Lu, W.; Li, Y.; Kinjo, R. Crystalline Tetraatomic Boron(0) Species. *J. Am. Chem. Soc.* **2019**, *141*, 5164–5168.
- (162) Masuda, J. D.; Schoeller, W. W.; Donnadiou, B.; Bertrand, G. Carbene Activation of P<sub>4</sub> and Subsequent Derivatization. *Angew. Chem., Int. Ed.* **2007**, *46*, 7052–7055.
- (163) Martin, C. D.; Weinstein, C. M.; Moore, C. E.; Rheingold, A. L.; Bertrand, G. Exploring the Reactivity of White Phosphorus with



Electrophilic Carbenes: Synthesis of a P<sub>4</sub> Cage and P<sub>8</sub> Clusters. *Chem. Commun.* **2013**, 49, 4486–4488.

(164) Masuda, J. D.; Schoeller, W. W.; Donnadieu, B.; Bertrand, G. NHC-Mediated Aggregation of P<sub>4</sub>: Isolation of a P<sub>12</sub> Cluster. *J. Am. Chem. Soc.* **2007**, 129, 14180–14181.

(165) Dorsey, C. L.; Squires, B. M.; Hudnall, T. W. Isolation of a Neutral P<sub>8</sub> Cluster by [2 + 2] Cycloaddition of a Diphosphene Facilitated by Carbene Activation of White Phosphorus. *Angew. Chem., Int. Ed.* **2013**, 52, 4462–4465.

## Recommended by ACS

### Stable Silapyramidanes

Taiki Imagawa, David Scheschkewitz, *et al.*

FEBRUARY 14, 2023

JOURNAL OF THE AMERICAN CHEMICAL SOCIETY

[READ](#) 

### Stabilized Molecular Diphosphorus Pentoxide, P<sub>2</sub>O<sub>5</sub>L<sub>2</sub> (L = N-Donor Base), in the Synthesis of Condensed Phosphate–Organic Molecule Conjugates

Kevin Qian, Christopher C. Cummins, *et al.*

MARCH 13, 2023

JOURNAL OF THE AMERICAN CHEMICAL SOCIETY

[READ](#) 

### An Air-Stable “Masked” Bis(imino)carbene: A Carbon-Based Dual Ambiphile

Ying Kai Loh, Guy Bertrand, *et al.*

JANUARY 17, 2023

JOURNAL OF THE AMERICAN CHEMICAL SOCIETY

[READ](#) 

### Synthesis, Isolation, and Characterization of Two Cationic Organobismuth(II) Pincer Complexes Relevant in Radical Redox Chemistry

Xiuxiu Yang, Josep Cornella, *et al.*

FEBRUARY 28, 2023

JOURNAL OF THE AMERICAN CHEMICAL SOCIETY

[READ](#) 

[Get More Suggestions >](#)